NATURAL RESOURCES

WASHINGTON GEOLOGY

VOL. 26, NO. 4
DECEMBER 1998

IN THIS ISSUE

WASHINGTON STATE DEPARTMENT OF
Natural Resources
Jennifer M. Belcher - Commissioner of Public Lands



WASHINGTON GEOLOGY

Vol. 26, No. 4
December 1998

Washington Geology (ISSN 1058-2134) is published four times each year by the Washington State Department of Natural Resources, Division of Geology and Earth Resources. This publication is free upon request. The Division also publishes bulletins, information circulars, reports of investigations, geologic maps, and open-file reports. A list of these publications will be sent upon request.

MAIN OFFICE
Department of Natural Resources
Division of Geology and Earth Resources
PO Box 47007
Olympia, WA 98504-7007

Phone: (360) 902-1450
Fax: (360) 902-1785

(See map on inside back cover for main office location.)

Internet Connections:
Library inquiries:
connie.manson@wadnr.gov
lee.walkling@wadnr.gov
Subscriptions/address changes:
geology@wadnr.gov

URL: http://www.wa.gov/dnr/htdocs/ger/ger.html

FIELD OFFICE
Department of Natural Resources
Division of Geology and Earth Resources
904 W. Riverside, Room 215
Spokane, WA 99201-1011

Phone: (509) 456-3255
Fax: (509) 456-6115
E-mail: robert.derkey@wadnr.gov

Cover Photo: Elaine Mustoe examines an unusual outcrop of metamorphic rocks uncovered during construction of a logging road in 1988. Located on Blanchard mountain, this outcrop of phyllite, green chert, and milky quartz contains enormous amounts of stilpnomelane, which has crystallized in a bed of Mesozoic chert cut by numerous quartz veins. The contact between metachert and phyllite can be seen at the center of the photo.

Crown Jewel Project and the Pend Oreille Mine—Status Report

Raymond Lasmanis, *State Geologist*
Washington Division of Geology and Earth Resources
PO Box 47007; Olympia, WA 98504-7007

The Crown Jewel Project covers a proposal by Battle Mountain Gold Company to mine by open-pit methods a large gold deposit on Buckhorn Mountain in Okanogan County. Recently some major milestones have been achieved to bring the mine towards production. On December 31, 1998, U.S. District Court of Oregon denied Okanogan Highland Alliance and the Colville Confederated Tribe's motion for a summary judgment. The Crown Jewel Final Environmental Impact Statement and Record of Decision were upheld. Subsequently, pursuant to Section 401 of the federal Clean Water Act, on January 14, 1999, the Washington Department of Ecology has approved a water-quality certification for the Crown Jewel mine. On January 20, 1999, Okanogan Superior Court Judge Burchard dismissed Okanogan Highlands Alliance's lawsuit against Okanogan County, Okanogan County Health District, and Battle Mountain Gold Company. The decision by the Okanogan Health District to defer solid waste permitting, which includes waste rock piles and mine tailings, to other required agency permits was upheld.

From 1906 to 1977, the Pend Oreille mine near Metaline Falls, Pend Oreille County, produced 14,796,305 tons of ore containing 166,985 tons of lead and 345,761 tons of zinc (Lasmanis, 1995). Cominco American, Inc., is proposing to reactivate the mine and mill to develop the Yellowhead ore zone. The Washington Department of Ecology is the lead agency and is in charge of conducting the Environmental Impact Statement (EIS). An environmental consulting firm hired by the Department of Ecology began work in August, 1998. The draft EIS is currently scheduled for completion in late summer, 1999.

Reference

Lasmanis, Raymond, 1995, History of the Metaline mining district and the Pend Oreille mine, Pend Oreille County, Washington: Washington Geology, v. 23, no. 1, p. 24-29. ■

Staff Notes

Kitty Reed, our editor of twelve years, left us in August to pursue personal interests. Kitty has been largely responsible for the high quality of Division publications. We will miss her a great deal and will try our best to maintain her high standards. She is a hard act to follow.

Carl Harris, our Senior Cartographer and GIS Specialist, was promoted to Computer Information Specialist 1 in November and now works for the Department of Natural Resources Information Technology Division sorting out the GIS hydro layer. We will miss Carl not only for his cartographic expertise, but also for his wry wit and willingness to serve as our guru of rock and mineral collecting.

Wendy Dixon-Shelton was hired in December to provide support for the front desk staff. She comes to us through Community Youth Services. We appreciate the work she is doing for us and the opportunity to provide a learning experience for her.

Stilpnomelane at Blanchard Mountain, Western Skagit County, Washington

George E. Mustoe
Geology Department
Western Washington University
Bellingham, WA 98225

Department of Natural Resources (DNR) forest lands in the southern Chuckanut mountains are a popular destination for hikers, horseback riders, and hang gliders, and since 1988, when an unusual outcrop of metamorphic rocks was uncovered during construction of a logging road, geologists and rockhounds as well. Located on Blanchard mountain, this outcrop of phyllite, green chert, and milky quartz contains enormous amounts of stilpnomelane (stilp-NOM-e-lane), a complex hydrous iron aluminosilicate that is usually found only as tiny crystals disseminated in iron-rich host rock. Few mineral enthusiasts have seen stilpnomelane under a microscope, let alone as the fist-size chunks that can be collected at this roadcut (cover photo, Fig 1).

I had the good luck to be the first geologist to visit the Blanchard mountain outcrop, but I was mystified by the sparkling black pea-sized crystal rosettes that comprise a major portion of the bedrock (Fig. 2). Western Washington University geology professor Edwin H. Brown had no difficulty identifying the specimens I collected as stilpnomelane, and it came as no surprise when x-ray diffraction data substantiated his diagnosis because Dr. Brown is one of the world's foremost authorities on this mineral. My own inability to identify the material is ironic, since thirty years ago my first geology-related job was helping construct the experimental apparatus Dr. Brown used in his studies of stilpnomelane geochemistry.

Despite its abundance of attractive mineral specimens, the Blanchard mountain site has received only brief printed mention (Gannaway, 1990). Stilpnomelane from Washington has previously been reported as a constituent of low-grade metamorphic rocks in the North Cascades (Brown, 1971). The mineral is common in rocks of the Shuksan Metamorphic Suite, particularly in the Finney Creek area where bedded magnetite contributed abundant amounts of iron to pore fluids during metamorphism. Despite having a similar chemical composition, Finney Creek stilpnomelane specimens are quite different in appearance from the material at Blanchard mountain (Fig. 3).

GEOLOGIC SETTING

The Chuckanut mountains are best known to geologists as the type locality for the Chuckanut Formation, a thick sequence of Early Tertiary arkosic sedimentary rock that underlies the northern part of the range. The southern Chuckanut region contains quite different bedrock—regionally metamorphosed Mesozoic marine sediments that have been tectonically transported from a distant location. The geology of this area is shown in Figure 4. Much of the bedrock is phyllite and fine-grained schist ('semischist') that belongs to the Shuksan Metamorphic Suite, an assemblage that originated when ancient ocean-floor sediments were subducted beneath the western edge of North America during the late Mesozoic. Elsewhere in the northwestern Cascades, the Shuksan Suite includes mafic schist that was produced by metamorphism of mid-ocean ridge basalt flows, but this rock type is absent in the Chuckanut region.

The southern Chuckanut mountains also contain other metamorphosed igneous rocks that have long puzzled geologists. These rocks form scenic outcrops at Windy Point, Pigeon Point, Bat caves, and Oyster dome. Gallagher and others (1988) believed that these meta-igneous materials represent pre-metamorphic compositional variations within the Shuksan Suite, but Whetten and others (1980) and Dragovich and others (1998) suggested they are remnants of some other thrust plate.

Stilpnomelane occurs within a zone of steeply dipping green chert that contains a complex pattern of hydrothermally deposited quartz veins. The chert unit is enclosed within a large body of phyllite. The vein-bearing zone is 150 m wide, with an exposed height of approximately 3 m. Stilpnomelane is particularly abundant in chert near the margins of quartz veins, and the crystalline veinlets and rosettes extend into the host rock for distances that vary from a few millimeters to 5 cm or

Figure 1. The stilpnomelane-bearing outcrop on the south flank of Blanchard mountain. *(See cover for a close-up view.)*

Figure 2. Rosettes of black stilpnomelane are abundant in chert bordering small veins of milky quartz.



Figure 3. A. Specimen from Blanchard mountain. Milky vein quartz contains stilpnomelane, but the mineral is not present in a 1 cm wide vein of translucent gray quartz. Cross-cutting relationships indicate that the gray vein post-dated stilpnomelane mineralization. **B.** Schist from Finney Creek contains stilpnomelane in two forms—veinlets composed of platy crystals and smaller crystal aggregates scattered throughout the surrounding matrix. Scale bar = 1 cm.

more. Stilpnomelane is also present as disseminated crystalline aggregates within small milky quartz veins; veins wider than a few centimeters contain little or no stilpnomelane.

MINERALOGY

Stilpnomelane was first described by Glocker (1827). The mineral has a layered crystal structure similar to that of talc or mica, being composed of a sheet of oxygen (O^{-2}) and hydroxyl (OH^{-}) ions and water molecules sandwiched between two silicate layers rich in iron and containing lesser amounts of aluminum (Eggleton, 1970, 1972; Deer and others, 1974). Individual triclinic crystals are seldom recognizable in hand specimens, and stilpnomelane typically appears as dark-colored micaceous flakes that are easily mistaken for biotite or chlorite.

Stilpnomelane characteristically forms stacks of thin plate-like crystals that are oriented radially at low angles of inclination around a central axis to produce an aggregate whose shape resembles a rose blossom that has been pressed between the pages of a book. In thin section, these masses are commonly visible as radiating clusters of needles or as bow-tie shaped sheaves. Both of these images are edge-on views of the sheet-like crystals (Fig. 5). The actual crystal arrangement is more accurately seen when uncut specimens are examined using a scanning electron microscope (Fig. 6).

The most important diagnostic characteristic is cleavage. The basal cleavage of stilpnomelane is less perfect than that of biotite or chlorite, yielding flakes that are brittle rather than elastic. Stilpnomelane may also be recognized by the presence of a weaker cleavage direction perpendicular to the basal plane. The optical properties of stilpnomelane are similar to those of biotite, with one important exception: stilpnomelane does not show the mottled extinction pattern that is observed in biotite. X-ray diffraction provides a reliable method of identification because the mineral produces a distinctive major peak near 12Å (Fig. 7). The position of this peak remains unchanged when the sample is treated with ethylene glycol or glycerin, in contrast to clay minerals that have 12Å lattice spacings (Chauvel, 1973).

CHEMICAL COMPOSITION

Stilpnomelane has a variable chemical composition that lies between two end members. Ferrostilpnomelane has an ideal formula of $K_5Fe^{2+}_{48}(Si_{63}Al_9)O_{168}(OH)_{216}\bullet 12H_2O$. The ideal chemical formula of the ferric end member is $K_5Fe^{3+}_{48}(Si_{63}Al_9)O_{216}\bullet 36H_2O$ (Eggleton and Chappell, 1978). Samples of both types also contain small amounts of sodium, calcium, and manganese.



Figure 4. Bedrock map of the southern Chuckanut mountains region based on mapping by Gallagher and others (1988).

Hutton (1938) applied the stilpnomelane name only to the ferric variety and referred to the oxidized form as ferristilpnomelane. Brown (1971) used 'green stilpnomelane' and 'brown stilpnomelane' to describe the ferrous and ferric varieties, but color is not always a reliable indicator of iron oxidation state. Ferrous stilpnomelane is typically olive green to pale green, but specimens that have a very low magnesium content may be brown. Oxidation typically causes green-colored stilpnomelane to become dark brown or reddish brown, but at Blanchard mountain and a few other locations, ferric stilpnomelane is an intense black. Chauvel (1973) suggested using the names ferrostilpnomelane and ferristilpnomelane. The stilpnomelane family also includes several varieties that contain large amounts of manganese substituted for part of the iron: ekmanite (Jakob, 1923), parsettensite (Deer and others, 1973; Guggenheim and Eggleton, 1994), and franklinphilite (Dunn and others, 1992).

The pure ferric and ferrous end members do not occur in nature, and individual specimens have compositions that lie somewhere between these extremes. Hashimoto (1969) observed that specimens tend to fall into either of two clusters based on their ferric/ferrous ratio. This bimodal distribution reflects the petrogenesis of stilpnomelane, where relatively pure ferrostilpnomelane may later become almost completely oxidized to ferristilpnomelane. This alteration trend is visible in thin section where green stilpnomelane grains are surrounded by brown rims (Hutton, 1938). Under laboratory conditions, Brown (1971) was able to reverse this reaction, producing green ferrostilpnomelane by heating brown ferristilpnomelane at 400 to 500°C for 27 days under pressures of 3,000 to 4,000 atmospheres.

The average chemical compositions of stilpnomelane from Blanchard mountain and Finney Creek are shown in Table 1. Water content is not reported for these samples because the



Figure 5. In thin section, stilpnomelane deceptively appears to consist of radial aggregates of needle-like crystals. Tiny clear zones within the radiating crystals are quartz inclusions. These Blanchard mountain specimens also show the fine microcrystalline structure of the chert matrix. Scale bars = 1 mm.

900°C temperature required for dehydration may cause the sample weight to change as a result of iron oxidation. Investigators have used special techniques to measure the iron content of stilpnomelanes from other locations, typically reporting H_2O^+ values of 5 to 9 percent (Chauvel, 1973).

The 49.57 percent SiO_2 content measured for the Blanchard mountain samples is somewhat greater than the 44 to 45 percent values that have been reported for most other stilpnomelanes. The Blanchard samples analyzed by the atomic absorption method probably have been contaminated by small amounts of quartz, visible in thin sections as tiny inclusions within the stilpnomelane crystals (Fig. 5). The 44.3 percent SiO_2 value that Dr. Foord obtained by electron microprobe analysis is closer to the 45.65 percent value determined by atomic absorption analysis for the quartz-free Finney Creek sample. For other elements, the Skagit County stilpnomelanes all fall within the compositional ranges that have previously been reported (Grüner, 1937; Hutton, 1945, 1956; Brown, 1971; Chauvel, 1973; Deer and others, 1974; Eggleton and Chappell, 1978).



Figure 6. This photo from a scanning electron microscope (SEM) of the broken surface of a Blanchard mountain specimen shows that stilpnomelane rosettes actually consist of flat plates that are inclined around a central axis. Although this is an accurate representation of the mineral's three-dimensional architecture, the SEM produces a black and white image based on electron emission that may be quite different from the specimen's appearance under visible light. In this example, the black stilpnomelane appears to be a light-colored mineral, and the quartz matrix is nearly black. This photograph demonstrates the SEM's great depth of field, allowing sharp focus even on a very irregular surface. Back-scattered electron image photographed using a Model 120 Cambridge Stereoscan electron microscope. Scale bar = 1 mm.



Figure 7. X-ray diffraction pattern obtained from Blanchard mountain stilpnomelane using Cu Kα radiation on a Rigaku Geigerflex diffractometer with graphite crystal monochromator. Finney Creek stilpnomelane yields a nearly identical x-ray pattern.

ORIGIN

Stilpnomelane is most commonly found in iron-rich rocks that have experienced greenschist- or blueschist-facies metamorphism as a result of exposure to relatively moderate temperatures and pressures. This mechanism accounts for the presence of the mineral in the North Cascades, the Franciscan Complex of the northern California Coast Range, in schist at Otago, New Zealand, and at numerous other sites around the world (Turner and Hutton, 1935). At Blanchard mountain, stilpnomelane

Table 1. Chemical analyses of Blanchard mountain and Finney Creek stilpnomelane. ND, no data

% oxide	Blanchard mtn.[1] (average of 3 analyses)	Blanchard mtn.[2] (average of 4 analyses)	Finney Creek[2] (average of 2 analyses)
SiO_2	44.3	49.57	45.65
Al_2O_3	6.25	5.91	7.57
TiO_2	0.0	0.0	0.0
MgO	1.2	1.31	3.99
CaO	ND	0.41	0.20
Na_2O	0.05	0.84	0.02
K_2O	1.2	0.93	1.17
MnO	2.15	2.23	4.36
Total iron as FeO	33.1	28.89	27.46
Total	88.25	90.09	90.42

[1] Electron microprobe data provided by Dr. Eugene Foord, U.S. Geological Survey, Denver, Colo., 1997.

[2] Atomic absorption spectrophotometry performed at Western Washington University using lithium metaborate fusion to dissolve samples.

crystallized during episodes of hydrothermal activity when iron was introduced into the silica-rich host rock by percolating water. The formation of stilpnomelane rather than some other iron mineral reflected physical and chemical conditions that were controlled by regional metamorphism.

Some Precambrian banded-iron formations contain stilpnomelane that formed during later metamorphism of the iron-rich sediment. Examples include the Lake Superior iron deposits of North America (Floran and Papike, 1975), the Hammersley Iron Range of Western Australia, and the Transvaal Supergroup of South Africa (Horstmann and Haelbich, 1995). Stilpnomelane also forms in certain metal sulfide deposits as a result of hydrothermal alteration of magnetite, pyrite, or pyrrhottite (Yui, 1962; Frondel, 1965).

In rare circumstances, ferristilpnomelane is produced by weathering of pyroxene. Eggleton (1975) described an example of this process at a skarn deposit near Canberra, Australia, where hedenbergite weathered to nontronite. This iron-rich variety of yellow smectite clay was in turn altered to brown ferristilpnomelane.

COLLECTING INFORMATION

The cluster of rounded summits east of Samish Bay is shown on maps and trail guides using two conflicting names: Blanchard Mountain and South Chuckanut Mountain. I have chosen to use the former name because it is consistent with the B prefix that the DNR uses to designate logging roads in this area. The stilpnomelane site is located on DNR property in the NW1/4 of sec. 14, T36 N, R3 E, of the Bow 7.5-minute quadrangle. The 1996 Blanchard Hill–South Chuckanut Mountain Trail Map published by the Pacific Northwest Trail Association is an excellent source of information about trails and logging roads (available for $4.95 from PNTA, 1361 Avon Allen Road, Mount Vernon, WA 28373).

To reach the outcrop from I-5, take Alger exit 240 and drive west toward Lake Samish. After 0.7 mi, turn left (south) on Barrel Springs Road and continue 0.9 mi. Turn right on DNR logging road B-1000, marked with a Blanchard Hill Trail sign. Follow this good gravel road for 1.9 mi. Just before reaching the upper trailhead for Lizard and Lily Lakes, turn left on road B-2000. Proceed 0.5 mi to reach the stilpnomelane-bearing outcrop, which is the second rock exposure beyond a culvert that crosses the headwaters of Whitehall Creek. The road continues an additional 2 mi to end at a spectacular viewpoint that overlooks Samish Bay and the San Juan Islands.

Stilpnomelane occurs at many locations in the North Cascade foothills near Finney Peak and Gee Point, south of the Cascade River near the town of Concrete. One site in the Mount Baker–Snoqualmie National Forest is very easy to reach. From Highway 20 just west of Concrete, cross the Skagit River on the Dalles Bridge. Continue east on South Skagit Highway for about 9 mi, then turn south on Forest Road 17 (Finney–Cumberland Road). Follow this road 10.5 mi to reach a large outcrop of thinly bedded siliceous magnetite bordered by dark-colored schist. Stilpnomelane is sparsely present in this schist as mica-like veinlets up to 1 cm in width and more abundant as disseminated microscopic crystals. The main geologic attraction of this site is the rich sedimentary iron ore, considered by early Skagit Valley settlers as a potential bonanza (Shedd and others, 1922; Zapffe, 1945).

ACKNOWLEDGMENTS

Thanks to Seattle mineral collector Robert O. Meyer for generously sharing electron microprobe data from analyses performed in 1997 by USGS mineralogist Eugene Foord. Northwest mineral collectors were saddened by Dr. Foord's death from cancer last year. My wife Elaine Mustoe served as a cheerful field companion and as an astute manuscript editor. E. H. Brown and J. D. Dragovich provided helpful reviews.

REFERENCES

Brown, E. H., 1971, Phase relations of biotite and stilpnomelane in the greenschist facies: Contributions to Mineralogy and Petrology, v. 31, no. 4, p. 275-299.

Brown, E. H., 1974, Comparison of the mineralogy and phase relations of blueschists from the North Cascades, Washington, and greenschists from Otago, New Zealand: Geological Society of America Bulletin, v. 85, no. 3, p. 333-344.

Chauvel, J. J., 1973, Les stilpnomelanes—Propriétés physiques et composition chemique: Contributions to Mineralogy and Petrology, v. 38, no. 1, p. 37-44.

Deer, W. A.; Howie, R. A.; Zussman, S., 1974, Rock-forming minerals: Longmans [London], p. 222-224.

Dragovich J. D.; Norman, D. K.; Grisamer, C. L.; Logan, R. L.; Anderson, Garth, 1998, Geologic map and interpreted geologic history of the Bow and Alger 7.5-minute quadrangles, western Skagit County, Washington: Washington Division of Geology and Earth Resources Open File Report 98-5, 80 p., 3 pl.

Dunn, P. J.; Peacor, D. R.; Su, S. C., 1992, Franklinphilite, the manganese analog of stilpnomelane: Mineralogical Record, v. 23, no. 6, p. 465-468.

Eggleton, R. A., 1970, Silicon tetrahedral sheet in stilpnomelane: Nature, v. 225, no. 5233, p. 625-626.

Eggleton, R. A., 1972, The crystal structure of stilpnomelane; Part II—The full cell: Mineralogical Magazine, v. 38, no. 298, p. 693-711.

Eggleton, R. A., 1975, Nontronite topotaxial after hedenbergite: American Mineralogist, v. 60, no. 11-12, p. 1063-1068.

Eggleton, R. A.; Chappell, B. W., 1978, The crystal structure of stilpnomelane; Part III—Chemistry and physical properties: Mineralogical Magazine, v. 42, no. 323, p. 361-368, M37.

Floran, R. J.; Papike, J. J., 1975, Petrology of the low-grade rocks of the Gunflint Iron-Formation, Ontario–Minnesota: Geological Society of America Bulletin, v. 86, no. 9, p. 1169-1190.

Frondel, Clifford; Ito, Jun, 1965, Stilpnomelane and spessartite-grossularite from Franklin, New Jersey: American Mineralogist, v. 50, no. 3-4, p. 498-501.

Gallagher, M. P.; Brown, E. H.; Walker, N. W., 1988, A new structural and tectonic interpretation of the western part of the Shuksan blueschist terrane, northwestern Washington: Geological Society of America Bulletin, v. 100, no. 9, p. 1415-1422.

Gannaway, W. L., 1990, Stilpnomelane of Blanchard mountain, Skagit County, Washington: Mineral News, v. 6, no. 4, p. 4.

Glocker, E. F., 1827, Beschreibung des Stilpnomelan, eines nuen schlesischen Fossils: Beyträge zur mineralogischen Kenntniss der Sudetenländer, inbesondere Schleisens [Breslau].

Guggenheim, S.; Eggleton, R. A., 1994, A comparison of the structures and geometric stabilities of stilpnomelane and parsettensite; A distance least-squares (DLS) study: American Mineralogist, v. 79, no. 5-6, p. 438-442.

Grüner, J. W., 1937, Composition and structure of stilpnomelane: American Mineralogist, v. 22, no. 8, p. 912-925.

Hashimoto, Mitsuo, 1969, A note on stilpnomelane mineralogy: Contributions to Mineralogy and Petrology, v. 23, no. 1, p. 86-88.

Horstmann, U. E.; Haelbich, I. W., 1995, Chemical composition of banded iron-formations of the Griqualand West Sequence, northern Cape Province, South Africa, in comparison with other Precambrian iron formations: Precambrian Research, v. 72, no. 1-2, p. 109-145.

Washington Mutual and SeaFirst Bank Win Project Impact Award

The first Project Impact Summit recognized those people and communities across the nation who have done outstanding work in disaster mitigation during the past year. The winners were announced during an awards banquet December 9 in Washington, D.C. Project Impact is a national initiative launched by the Federal Emergency Management Agency (FEMA) to include individuals, government officials, and business leaders in a partnership to make their communities resistant to disasters.

FEMA Director James Lee Witt said, "The people of every Project Impact community have made a commitment to identify and prioritize disaster risk....These awards are to honor the hard work and dedication of our partners nationwide."

The Model National Corporate Partners Award went to Washington Mutual and SeaFirst Bank/Bank America in Seattle, Wash., for their expanded damage-prevention efforts against the risk of Pacific Northwest earthquakes. Teaming up with insurance companies like SAFECO and PEMCO and building professional associations, they are actively leading the development of a new Home Earthquake Retrofit Program, which is well on its way to retrofitting 2,000 homes.

Project Impact has also published a Hazard Mitigation Guidebook for Northwest Communities—Alaska, Idaho, Oregon, Washington. The guidebook gives practical measures to prepare for disasters such as floods, landslides, earthquakes, tsunamis, wildfires, windstorms, blizzards, and volcanic eruptions. For more information or copies of the guidebook, contact FEMA; Federal Regional Center, Region 10; 130 228th St SW; Bothell, WA 98021-9796; or (425) 487-4678.

Hutton, C. O., 1938, The stilpnomelane group of minerals: Mineralogical Magazine, v. 25, no. 163, p. 172-206.

Hutton, C. O., 1945, Additional optical and chemical data on the stilpnomelane group of minerals: American Mineralogist, v. 30, no. 11-12, p. 714-718.

Hutton, C. O., 1956, Further data on the stilpnomelane mineral group: American Mineralogist, v. 41, no. 7-8, p. 608-615.

Jakob, J., 1923, Vier Mangansilikat aus dem Val d'Err (Kt Graubunden): Schweizerische Mineralogische und Petrographische Mitteilungen, v. 3, p. 227-237.

Shedd, Solon; Jenkins, O. P.; Cooper, H. H., 1922, Iron ores, fuels and fluxes of Washington: Washington Division of Geology Bulletin 27, 160 p., 1 pl.

Turner, F. J.; Hutton, C. O., 1935, Stilpnomelane and related minerals as constituents of schists from western Otago, New Zealand : Geological Magazine, v. 72, no. 847, p. 1-8.

Whetten, J. T.; Zartman, R. E.; Blakely, R. J.; Jones, D. L., 1980, Allochthonous Jurassic ophiolite in northwest Washington: Geological Society of America Bulletin, v. 91, no. 6, p. 359-368.

Yui, S., 1962, Stilpnomelane from the Motoyasu Mine, Shikoku: Journal of the Geological Society of Japan, v. 68, p. 597-600.

Zapffe, Carl, 1945, Hamilton iron-bearing lenses on Iron Mountain, Skagit County, Washington: Mining World, v. 7, no. 11, p. 25-29. ■

Reidel and Gephart Win Fitzner/Eberhardt Award

Steve Reidel and Roy Gephart of Battelle's Applied Geology and Geochemistry Group in Richland, Wash., were recently named recipients of the 1998 Pacific Northwest National Laboratory's Fitzner/Eberhardt Award for Outstanding Contributions to Science and Engineering Education.

Dr. Reidel has mentored many college students at PNNL through the Association of Western Universities Program and is an adjunct professor of geology at Washington State University. Steve also teaches introductory courses in astronomy and geology at Yakima Community College and helps local K-12 teachers improve their knowledge of geology. Steve writes a periodic column in the Tri-City Herald featuring regional geology and gives many public lectures each year to groups ranging from clubs to professional societies to school classrooms. Steve has about 100 publications (articles, maps and abstracts). He was coauthor with Karl Fecht of our Open File Reports 94-8 and 94-13. He also acted as a supplier of geologic data, leader of field trips, and reviewer of texts for our 1:100,000 and 1:250,000 (GM-45) maps of southeastern Washington.

Roy Gephart is a geohydrologist with 25 years' experience in the hazardous waste industry. He frequently works with local and national news media on astronomy, geohydrology, and Hanford waste generation and cleanup and gives science presentations in community forums and to students (elementary through graduate level). Roy formed the Tri-City Astronomy Club and organizes community star gazes. He is a board member of a nonprofit organization renovating the Rattlesnake Mountain Observatory for online student education, teacher training, and research use. Roy has authored more than 50 publications, including textbooks and science-related articles/photographs in national magazines. In 1996, he wrote an award-winning publication on the history of Hanford's high-level radioactive tank waste. He is now working on a new publication summarizing the history of waste generation/release at Hanford and the decision-making challenges of site cleanup.

CD-ROM REVIEW: The Photo-Atlas of Minerals—A comprehensive reference tool for mineral enthusiasts, students, teachers, and professional mineralogists

produced by The Gem & Mineral Council, 1998
Los Angeles County Museum of Natural History
900 Exposition Blvd, Los Angeles, CA 90007
$49.95 plus $5.00 for shipping & handling

The Photo-Atlas of Minerals was developed by Dr. Anthony R. Kampf, curator of Mineral Sciences at the Los Angeles County Museum, and Dr. George Gerhold of Western Washington University. It contains more than 6,500 high-resolution images by well-known mineral photographers, among them Dr. Wendell E. Wilson, Louis Perloff, and Rick Dillhoff of Washington State. More than 6,400 of the images are in color; the others are scanning electron microscope (SEM) photos. The mineral property data were compiled by Lanny Ream.

More than 800 different mineral species are illustrated. Descriptive data on these and thousands of other minerals presents coverage of the entire mineral kingdom. Other features include an audio-based mineral pronunciation guide, a linked glossary of mineral terms, the latest mineral classification scheme according to Strunz, and cross-indexing capabilities for easy searches. System requirements are: a PC 486-33 or better, CD-ROM drive, sound card, Windows 3.1, 95, 98, or NT, a mouse, and for extra sharp images a 24-bit graphics card.

The Pacific Northwest is well represented. For Washington State, 61 mineral species are illustrated from 36 localities. Of these, there are multiple illustrations, such as 31 photos of quartz, 21 of pyrite, and 9 of gold specimens. Oregon is represented by 32 localities with 34 species, mostly zeolites. Milton Speckels is credited with much of the Oregon photography. Idaho is represented by 8 localities and 13 species, including 40 photos of various pyromorphite crystal groups. Montana has 8 localities with 23 species and British Columbia has 8 localities and 14 species.

A number of years ago, I was involved in the early planning of this product with Dr. George Gerhold. It is now available and has given me many hours of enjoyment. Being the first of its kind, there are bound to be a few minor glitches, considering the volume of data. For instance, pink zoisite from Tunk Creek, Okanogan County, is mislabeled as coming from Pacific County, and I am still trying to find out where one would find caledonite at Chehalis, Lewis County.

The first release of the Photo-Atlas has sold out. The upgrade (version 1.1) incorporates a number of improvements and fixes most of the glitches in the previous version. My comments on the locations of the pink zoisite and caledonite came too late to make it into this upgrade. Dr. Kampf promises to fix these and any other glitches in the next version and asks that users let them know if anything further turns up.

The most significant enhancements are a much more detailed hierarchical implementation of the Strunz mineral classification system and full implementation of the 'Combine Indexes' feature, making it a more powerful database searching tool. No new minerals or images have been added. Users who purchased version 1.0 may upgrade to version 1.1 for $9.95. Further information regarding the Photo-Atlas of Minerals and an order form are online at the Gem and Mineral Council webpage at http://nhm.org/~gmc.

The Photo-Atlas of Minerals is reasonably priced and comes very highly recommended.

Raymond Lasmanis



Barite
Spruce claim, King Co., Washington
Owner: Rick Dillhoff
Photographer: Rick Dillhoff

SEM Photo

Okanoganite on Euxenite
Washington Pass, Okanogan Co., Washington
Owner: Donald Howard
Photographer: Donald Howard

SEM Photo

Zircon
Washington Pass, Okanogan Co., Washington
Owner: Donald Howard
Photographer: Donald Howard

Sample illustrations of minerals from Washington State. The barite photo is in color on the CD-ROM. The other two photos are SEM images. An SEM image is an accurate representation of a mineral's three-dimensional architecture, but since the SEM produces a black and white image based on electron emission, it may be quite different from the specimen's appearance under visible light.

Washington State Gem & Mineral Clubs

The following list of gem and mineral clubs and organizations in Washington State is as up-to-date as we could make it. To make any additions or corrections, contact our front desk by *phone* (360)902-1450, *fax* (360)902-1785, or *e-mail* geology@wadnr.gov.

ABERDEEN

Chehalis Valley Gem Club
422 W 2nd St; Aberdeen, WA 98520
Robert Musgrove, President
Meetings: Second Thursday, 1:00 pm,
in members' homes

Grays Harbor Geology & Gem Society
PO Box 2003; Aberdeen, WA 98520
Leonard Airhart (360)533-3078
Vaughn Hamilton, President
Meetings: Third Friday, 7:30 pm,
Pearsall Multi-Purpose Center
2109 Summer Ave.; Aberdeen, WA

ARLINGTON

Port Susan Gem & Mineral Club
9406 16th St NE;
Arlington, WA 98223-8904
Meetings: First Wednesday, 12:00 pm,
12015 Marine Dr; Marysville, WA

BELLEVUE

Bellevue Rock Club, Inc.
Box 1851; Bellevue, WA 98009-1851
Elanor Dickson (425)746-8412
Odella Jackson, President
Meetings: First Tuesday, 7:00 pm,
Hyaki School,
445 128th Ave SE; Bellevue, WA

BELLINGHAM

Mt. Baker Rock & Gem Club
PO Box 142; Bellingham, WA 98227
Homer Owens (360)647-6699
Rosemae Bork, President
Meetings: Third Monday, 7:30 pm
Bloedel Donovan Park Pavilion

BREWSTER

Chief Joseph Gem & Mineral Club
PO Box 51; Brewster, WA 98812
Meetings: Second Wednesday, 7:30 pm
Shulls Towing & Wrecking Yard Office

CHELAN

Lake Chelan Rock & Mineral Club
PO Box 487; Manson, WA 98831
Hal Porter (509)687-9615
Charles Leffler, President
Meetings: First Tuesday, 7:00 pm,
Lake Chelan Recl. Dist. Board Room

COLLEGE PLACE

Blue Mountain Gem & Mineral Society
1011 Broadway;
College Place, WA 99324-1571
(509)525-1776
Pearl Vickroy, President
Meetings: Second Thursday, 7:30 pm
Rt 1 Box 114, Walla Walla, WA

EDMONDS

Maplewood Rock & Gem Community Club
PO Box 1295; Edmonds, WA 98133
21318 Pioneer Way; Edmonds, WA 98020
Jeanette Koop (425)771-4687
Marcy Kleckner, President
Meetings: Third Monday, 8:00 pm
Club House, 8802 196th St SW

ELK

Pend Oreille Rock & Gem Club
42503 Regal Rd; Elk, WA 99009
Lois or Bill Hunter (509)292-2485
Bill Hunter, President
Meetings: Third Thursday, 6:30 pm
Fertile Valley Grange

EVERETT

Everett Rock & Gem Club, Inc.
PO Box 1615; Everett, WA 98206
George Burkhart, President (425)672-8952
Fritz Mack, Vice President (425)513-0115
Meetings: Second Monday, 7:30 pm
Our Savior Lutheran Church
215 Mukilteo Blvd; Everett, WA

FEDERAL WAY

Federal Way Gem Club
PO Box 3305;
Federal Way, WA 98063-3305
(253)661-0746
Keith Ray (206)854-1303
Carol Uhlman, President
Meetings: Third Friday, 7:30 pm
Totem Jr. High School Library

FORT LEWIS

Fort Lewis Rock Club
10107 111th St Ct SW; Tacoma, WA 98498
Meetings: First Monday, 7:00 pm
MWR Arts & Crafts Center, Bldg. #5038

KENNEWICK

Lakeside Gem & Mineral Club
PO Box 6652; Kennewick, WA 99336-9998
(509)783-4262/(509)967-3138
Larry Hulstrom, President
Meetings: First Wednesday, 7:00 pm
First Savings Bank of Washington,
203 W 1st Ave

Rockateers Gem & Mineral Club
(Junior Club)
PO Box 6652; Kennewick, WA 99336
Meetings: Second Tuesday, 6:30 pm
First Savings Bank of Washington,
basement, 203 W 1st Ave

LANGLEY

Whidbey Pebble Pushers
PO Box 279; Langley, WA 98260
Gen Richards (360)221-2637
Charles Bash, President
Meetings: First Wednesday, 7:00 pm
4-H Club Bldg, Island Co. Fairgrounds

LONGVIEW

Southern Washington Mineralogical Society
PO Box 704; Longview, WA 98632-7451
Emmet Johnson, President (360)636-3491
Meetings: Fourth Saturday, 7:00 pm
Catlin Grange Hall;
207 Shawnee St N, Kelso, WA

MARYSVILLE

Marysville Rock & Gem Club, Inc.
4406 92nd St NE; Marysville,
WA 98270-2506
Lloyd Bellman (360)659-2554
Ed Lehman (206)334-6282
Dale Sanders, President
Meetings: Second Tuesday, 7:30 pm
Jennings Park Barn

MOUNT VERNON

Northwest Rockies Family 4-H Club
(Junior Club)
6131 Dale Way; Lynnwood, WA 98036
(253)856-1564/(206)776-6598
Aaron Bever, President
Meetings: Second Monday, 7:00 pm
Upstairs Skagit Co. Fair Office

Skagit Rock & Gem Club
PO Box 244; Mount Vernon, WA 98273
Dave Britten (360)755-0741
Meetings: First Wednesday, 7:30 pm
IBEW Hall; 706 W Division

OAK HARBOR

Whidbey Island Gem Club
PO Box 224; Oak Harbor, WA 98277
W. H. Schreiter (360)678-5485
Keith Ludeman, President
Meetings: Second Wednesday, 7:00 pm
Oak Harbor Senior Center

OLYMPIA

Washington Agate & Mineral Society
PO Box 2553; Olympia, WA 98507
M. J. Huetter (360)459-8121
Curtis Mack, President
Meetings: First Tuesday, 7:00 pm
First Baptist Church; 22nd & College,
Lacey

PORT ANGELES

Clallam County Gem & Mineral Society
PO Box 2624; Sequim, WA 98382
John Schuy, President
Meetings: Third Wednesday, 7:00
Sequim Senior Center
921 E Hammond St, Sequim, WA

PORT ORCHARD

Kitsap Mineral & Gem Society
PO Box 3462; Silverdale, WA 98383
(360)697-1859
Jim McClure (253)265-3011
Mark Stephens, President
Meetings: Second Friday, 7:30 pm
Chico Alliance Church Gym,
3670 Chico Way; Bremerton, WA
e-mail: pogy2@worldnet.alt.net

PORT TOWNSEND

Port Townsend Rock Club
8275 Hwy 20; Port Townsend, WA 98368
Marie Anderson (360)385-0420
Janet Smith, President
Meetings: Last Monday, 7:30 pm
Jefferson County Fairgrounds

PUYALLUP

Puyallup Valley Gem & Mineral Club
PO Box 134; Puyallup, WA 98371-0014
Steve Dugan (206)531-2484
Gene Beckstead (253)535-6536
Ralph Graves, President
Meetings: Second & Fourth Friday, 7:30 pm
Fruitland Grange Hall,
112th St E & 87 Ave E

REDMOND

East Kingco Rock Club, Inc.
PO Box 2203; Redmond, WA 98073-2203
Jack Frasl (206)820-0244
Jack Donner, President
Meetings: Fourth Monday, 7:30 pm;
(Juniors 7:00 pm)
Rose Hill Presbyterian Church
Fellowship Hall

SEATTLE

Boeing Employees Mineralogical Society
The Boeing Co., Box 3707 MS-8L-35;
Seattle, WA 98124-2207
Lee Adams, President (206)235-1338
Meetings: Second Thursday, 7:30 pm
Boeing Activities Center, Rm B
22649 83rd Ave S, Kent, WA

Issaquah Valley Rock Club, Inc.
1706 NW 59th, #303; Issaquah, WA 98107
(206)789-0588
Mike S. Tanaka, President (425)392-7858
Meetings: Last Friday, 7:30 pm
Issaquah Community Center
180 E Sunset Way

North Seattle Lapidary & Mineral Club, Inc.
(206)524-5188
Susan Gardner (206)782-6522
Harold Cutton, President
Meetings: Second Tuesday, 8:00 pm
(Juniors 7:00)
Northminster Presbyterian Church
7706 25th Ave NW, basement of church

Seattle Faceting Club
Leonard Bahr (206)242-5560
Norman Steele, President
Meetings: Fourth Tuesday, 7:30 pm,
in members' homes

South Seattle & Des Moines
Gem & Mineral Club
13056 24th Ave. S; Seattle, WA 98168
Bill Scott, President
Meetings: Third Thursday, 7:00 pm
Tukwila Senior Center, 4101 S 131st St

West Seattle Rock Club
PO Box 16145; Seattle, WA 98116
Paul Schoeler (206)932-1522
Meetings: Fourth Wednesday, 7:30 pm
(Except July and December;
November, Third Wednesday)
Adams Hall,
Tibbetts United Methodist Church,
3940 41st St SW, Seattle, WA
e-mail: tobycoz@aol.com

SHELTON

Shelton Rock & Mineral Society
PO Box 242; Shelton, WA 98584
(360)427-0387
Bill Barron, President
Meetings: Last Thursday, 7:00 pm
Mason Count PUD #3 Bldg., 307 W Cota

SNOHOMISH

Snohomish Lapidary Club
9108 105th Ave SE;
Lake Stevens, WA 98258-8919
Don Brown (360)659-6444
Glenn Morita (425)337-0385
Larry Soderblom, President
Meetings: First Monday, 7:30 pm
Snohomish Public Library, First & Cedar
e-mail: gmorita@seanet.com

SPOKANE

Columbia Geological Society, Inc.
W 1517 Carlisle Ave;
Spokane, WA 99205-3511
(509)328-6584
Jean Williams, President
Meetings: First Thursday, 7:30 pm
(except July, August, and December)
Manito Garden Center, Manito Park

Rock Rollers Club, Inc.
OPP Station, PO Box 14766;
Spokane, WA 99214-0766
Mable Rutherford (509)926-6851
Ed Brandstoettner (509)467-0360
Leon Agee, President
Meetings: Fourth Saturday, 7:30 pm
East Spokane Grange Hall; N 1621 Park Rd.

TACOMA

Tacoma Faceters Guild
5109 Point Fosdick Dr NW;
Gig Harbor, WA 98335
Meetings: Fourth Monday, 7:00 pm
Fircrest Community Center,
555 Contra Costa; Tacoma, WA

TENINO

Tenino Rock Cruisers
Box 4008; Tenino, WA 98589
Ken Hedden, President 264-2570
Meetings: Second Wednesday, 7:00 pm
(except August and December)
Tenino Masonic Hall, 260 W Sussex

TUKWILA

Northwest Opal Association
Meetings: Second Wednesday
(except July and August)
Tukwila Denny's; 5700 Southcenter Blvd
(north side of I-405)

WALLA WALLA

Marcus Whitman Gem &
Mineral Society, Inc.
PO Box 338;
Walla Walla, WA 99362-0009
(509)529-3673
Jack Edwards, President
Meetings: Second Tuesday, 7:30 pm
Lions Park Field House,
Larch St, College Place

WASHOUGAL

Washougal Gem Club, Inc.
938 17th St; Washougal, WA 98671-1510
Dolphe S. Gilbreath, President
Meetings: Fourth Tuesday, 2:00 pm
(except July & August), 938 17th St.

WENATCHEE

Gingko Mineral Society
PO Box 303; Wenatchee, WA 98807
Claude Johnston (509)884-3188
Mike Edgett, President
Meetings: Second & Fourth Friday, 7:30 pm
Memorial Hall; 215 Okanogan Ave

WOODINVILLE

Maplewood Rock & Gem Club
20718 59th Ave SE;
Woodinville, WA 98072
Bud Lischke (206)365-5312
Meetings: Third Monday
8802 196th St SW, Edmonds, WA

YAKIMA

Yakima Gem & Mineral Club
PO Box 969; Yakima, WA 98907
Andy Beeman (509)457-6339
Bill Snell (509)837-5329
Jack Friedt, President
Meetings: Third Friday, 7:30 pm
Central Lutheran Church;
16th & Yakima Ave

YELM

Nisqually Valley Rockhounds Society
PO Box 561; Yelm, WA 98597
(206)491-1429
Tim Howard, President
Meetings: Second Thursday, 7:30 pm
Yelm High School Art Room

STATE AND NATIONAL

Northwest Federation of Mineralogical
Societies (NFMS)
4401 SW Hill St; Seattle, WA 98116
(206)937-7872
Dorothy Lee, President

Northwest Micro Mineral Study Group
(NWMMSG)
Donald G. Howard
356 SE 44th Ave
Portland, OR 97215

Friends of Minerology, Inc.
Pacific Northwest Chapter
Bob Mayer, President (425)641-0723
John Cornish (360)457-7630
16239 NE 18th St; Bellevue, WA 98008

Washington State Mineral Council
Bob O'Brien, President
1665 S Elger Bay Rd;
Stanwood, WA 98292

Selected Additions to the Library of the Division of Geology and Earth Resources

August 1998 through January 1999

THESES

Bacon, D. H., 1997, Residence time of labile carbon in the vadose zone: Washington State University Doctor of Philosophy thesis, 256 p.

Christenson, D. H., 1997, Hydrology of urban watersheds in Cheney, Washington with an emphasis on stormwater runoff and water quality: Eastern Washington University Master of Science thesis, 139 p.

Fiedorowicz, B. K., 1997, Geologic evidence of historic and prehistoric tsunami inundation at Seaside, Oregon: Portland State University Master of Science thesis, 197 p.

García, A. F., 1996, Active tectonic deformation and late Pleistocene and Holocene geomorphic and soil profile evolution in the Dosewallips River drainage basin, Olympic Mountains, western Washington State: University of New Mexico Master of Science thesis, 152 p., 4 plates.

Kirn, S. L., 1995, Petrology of a Mount Rainier pyroclastic eruption: Brown University Bachelor of Science thesis, 1 v.

Martin, W. J., 1996, Integration of risk analysis and sorption studies in the subsurface transport of aqueous carbon-14 at the Hanford site: Washington State University Doctor of Philosophy thesis, 201 p.

Includes:

Martin, W. J.; Whelan, Gene; Conca, J. L., 1996, Integration of sorption and risk assessment in waste disposal. p. 30-50.

Martin, W. J.; Whelan, Gene, 1996, Modeling of radioactive transport from decommissioned nuclear reactor waste. p. 6-29.

Massong, T. M., 1998, Influence of lithology, sediment supply, and log jams on the distribution of bedrock and alluvial channels: University of Washington Master of Science thesis, 62 p.

Matthews, J. M., 1996, The stratigraphy and sedimentology of the Eocene O'Brien Creek Formation, Okanogan Highlands, north-east Washington, USA: Washington State University Doctor of Philosophy thesis, 182 p.

McArdell, B. W., 1997, Field experiments on the controls of downstream fining in gravel-bed rivers: Johns Hopkins University Doctor of Philosophy thesis, 137 p.

McNeill, L. C., 1998, Structure and seismic hazards of the offshore Cascadia forearc and evolution of the Neogene forearc basin: Oregon State University Doctor of Philosophy thesis, 178 p., 1 plate.

Includes:

McNeill, L. C.; Goldfinger, Chris; Hummon, Cheryl, 1998, Neotectonic map of the Washington and northwest Oregon continental margin. 1 plate.

Meng, Xiangying, 1997, Radiolarian biostratigraphy of the Upper Jurassic of San Pedro del Gallo terrane, north-central Mexico, and the Lower Cretaceous of Nooksack Group, Nooksack terrane, northwestern Washington: University of Texas at Dallas Doctor of Philosophy thesis, 351 p.

Olson, P. L., 1995, Shallow subsurface flow systems in a montane terrace–floodplain landscape—Sauk River, North Cascades, Washington: University of Washington Doctor of Philosophy thesis, 277 p.

Payne, C. W., 1998, Lithofacies, stratigraphy, and geology of the middle Eocene type Cowlitz Formation and associated volcanic and sedimentary units, eastern Willapa Hills, southwest Washington: Oregon State University Master of Science thesis, 253 p., 3 plates.

Phillips, W. M., 1997, Applications of noble gas cosmogenic nuclides to geomorphology: University of Arizona Doctor of Philosophy thesis, 256 p.

Silva de Echols, Catarina, 1995, Epibionts and their effects on the taphonomy of Recent crinoid ossicles, Friday Harbor, Washington, and San Salvador, Bahamas: Auburn University Master of Science thesis, 161 p.

Wills, M. T., 1998, Dairy farming and the effects of agricultural, nonpoint-source pollution on stream water quality, Johnson Creek watershed, Whatcom County, Washington: Western Washington University Master of Science thesis, 282 p.

U.S. GEOLOGICAL SURVEY

Published Reports

Page, R. A.; Basham, P. W., 1985, Earthquake hazards in the offshore environment: U.S. Geological Survey Bulletin 1630, 69 p.

Solley, W. B.; Pierce, R. R.; Perlman, H. A., 1998, Estimated use of water in the United States in 1995: U.S. Geological Survey Circular 1200, 71 p.

Thompson, R. S.; Hostetler, S. W.; Bartlein, P. J.; Anderson, K. H., 1998, A strategy for assessing potential future changes in climate, hydrology, and vegetation in the western United States: U.S. Geological Survey Circular 1153, 20 p.

Todd, V. R., 1995, Geologic map of the Doe Mountain 15′ quadrangle, Okanogan County, Washington: U.S. Geological Survey Miscellaneous Field Studies Map 2306, 1 sheet, scale 1:62,500, with 17 p. text.

U.S. Geological Survey, 1998, Mineral industry surveys—Washington: U.S. Geological Survey, 7 p.

Fact Sheets, Open-File, and Water-Resources Investigations Reports

Derkey, P. D.; Johnson, B. R.; Lackaff, B. B.; Derkey, R. E., 1998, Digital geologic map of the Rosalia 1:100,000 quadrangle, Washington and Idaho—A digital database for the 1990 S. Z. Waggoner map: U.S. Geological Survey Open-File Report 98-357, 27 p.

Drost, B. W.; Turney, G. L.; Dion, N. P.; Jones, M. A., 1998, Hydrology and quality of ground water in northern Thurston County, Washington: U.S. Geological Survey Water-Resources Investigations Report 92-4109 (revised), 230 p., 6 plates.

Embrey, S. S., 1991, Available habitat for salmon and steelhead trout in the lower Puyallup, White, and Carbon Rivers in western Washington: U.S. Geological Survey Water-Resources Investigations Report 89-4125, 62 p.

Finn, C. A.; Brenner, K. C.; McCafferty, Anne; Kucks, R., 1998, Washington aeromagnetic maps and data: U.S. Geological Survey Open-File Report 98-241, 1 v. [downloaded 9/30/98 from http://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-98-0241/wash.html]

Harp, E. L.; Chleborad, A. F.; Schuster, R. L.; Cannon, S. H.; Reid, M. E.; Wilson, R. C., 1998, Landslides and landslide hazards in Washington State due to February 5–9, 1996 storm: U.S. Geological Survey Administrative Report, 1 v. [downloaded 9/25/98 from http://geohazards.cr.usgs.gov/html_files/nlic/Wash_hrp/wash1.htm]

Hoblitt, R. P.; Walder, J. S.; Driedger, C. L.; Scott, K. M.; Pringle, P. T.; Vallance, J. W., 1998, Volcano hazards from Mount Rainier, Washington, revised 1998: U.S. Geological Survey Open-File Report 98-428, 11 p., 2 plates.

Inkpen, E. L.; Embrey, S. S., 1998, Nutrient transport in the major rivers and streams of the Puget Sound basin, Washington: U.S. Geological Survey Fact Sheet 009-98, 4 p.

Johnson, B. R.; Derkey, P. D.; Frost, T. P.; Derkey, R. E.; Lackaff, B. B., 1998, Digital geologic map of Spokane County and vicinity, Washington and Idaho: U.S. Geological Survey Open-File Report 98-503, 42 p.

Kahle, S. C., 1998, Hydrogeology of Naval Submarine Base Bangor and vicinity, Kitsap County, Washington: U.S. Geological Survey Water-Resources Investigations Report 97-4060, 107 p., 7 plates.

Kresch, D. L., 1998, Determination of upstream boundaries on western Washington streams and rivers under the requirements of the Shoreline Management Act of 1971: U.S. Geological Survey Water-Resources Investigations Report 96-4208, 57 p., 4 plates.

Kresch, D. L., 1998, Determination of upstream boundary points on northeastern Washington streams and rivers under the requirements of the Shoreline Management Act of 1971: U.S. Geological Survey Water-Resources Investigations Report 98-4160, 19 p.

LaHusen, R. G., 1998, Detecting debris flows using ground vibrations: U.S. Geological Survey Fact Sheet 236-96, 2 p.

Mastin, M. C., 1998, Flood potential of South Prairie Creek, Pierce County, Washington: U.S. Geological Survey Water-Resources Investigations Report 98-4009, 75 p.

Melick, Roger; Furman, T. L.; Langer, W. H.; van Brenk, H., 1998, Using geographic information systems with specific examples for the aggregate industry: U.S. Geological Survey Open-File Report 98-39, 1 v. [downloaded from http://minerals.cr.usgs.gov/publications/ofr/98-39/report-.htm]

Scott, K. M.; Wolfe, E. W.; Driedger, C. L., 1998, Mount Rainier—Living with perilous beauty: U.S. Geological Survey Open File Report 98-428, Fact Sheet 065-97, 4 p.

Soller, D. R., editor, 1998, Digital mapping techniques '98—Workshop proceedings: U.S. Geological Survey Open-File Report 98-487, 134 p.

Sumioka, S. S., 1998, A survey of ground-water quality in the Toppenish Creek Basin, Yakama Indian Reservation, Washington, 1989–91: U.S. Geological Survey Water-Resources Investigations Report 97-4194, 89 p., 5 plates.

U.S. Geological Survey, 1996, Annual project summaries—Volume 37, Pacific Northwest: U.S. Geological Survey, 115 p. [reports downloaded Dec. 2, 1998, from http://erp-web.er.usgs.gov/reports/VOL37/pn_vol37.htm]

Includes:

Atkinson, G. M.; Boore, D. M., 1996, Ground motion relations for the Pacific Northwest. p. 1-2.

Atwater, B. F., 1996, Prehistoric earthquakes in western Washington. p. 3-4.

Brocher, T. M., 1998, Analysis of wide-angle seismic data from Cape Blanco, Oregon. p. 22-24.

Carpenter, J. E., 1996, Workshops to explain the changes resulting from seismic rezone from UBC Zone 2B to Zone 3 in southwest Washington State. p. 15-17.

Creager, K. C.; Malone, S. D., 1996, Broad-band array analysis of the Puget Sound region. p. 18-20.

Crosson, R. S., 1996, Earthquake hazard research in the Pacific Northwest using Washington Regional Seismograph Network data. p. 21-24.

Dadisman, S. V.; Johnson, S. Y.; Childs, J. R., 1996, Neotectonics of the northern Puget Lowland and eastern Straits of Juan de Fuca, Washington. p. 25.

Doser, D. I., 1998, Source processes of crustal earthquakes and their relation to the tectonics of southern Alaska. p. 55-57.

Endo, E. T.; Dzurisin, Daniel; Iwatsubo, E. Y., 1996, Physical processes in large silicic magma systems. p. 35-36.

Fleming, S. W.; Trehu, A. M., 1996, Configuration of the subduction zone backstop and its influence on forearc deformation—Constraints from modeling gravity and magnetic anomalies. p. 99-109.

Graves, R. W., 1996, Simulating the 3D basin response in the Portland and Puget Sound regions from large subduction zone earthquakes. p. 38-44.

Hyndman, R. D.; Wang, Kelin; Dragert, Herb, 1996, Cascadia megathrust earthquake potential—Constraints from current deformation and the thermal regime. p. 49-60.

Jacoby, G. C., 1996, Tree-ring dating of coseismic coastal subsidence in the Pacific Northwest region. p. 61-64.

Johnson, S. Y., 1996, Cenozoic tectonic and paleogeographic evolution, Puget Lowland seismic zone, Washington. p. 65-67.

Kelsey, H. M.; Witter, R. C., 1998, A relative sea level history and an assessment of active upper-plate faulting, Coquille River estuary, Oregon. p. 64-70.

Kulm, L. D.; Goldfinger, Chris; Yeats, R. S., 1996, Cascadia subduction zone—Neotectonics of the continental shelf off Oregon and Washington. p. 68-74.

Mader, G. G., 1998, Preparation of a guide to using relative earthquake hazard maps for mitigating seismic risk in the Portland metropolitan area—Collaborative research with Metro and Spangle Associates. p. 94-95.

Madin, I. P.; Wang, M. M., 1998, Earthquake hazard mapping for Oregon communities. p. 84.

Malone, S. D., 1996, Seismic velocity structure of the greater Mount Rainier area. p. 76-78.

McCalpin, J. P., 1996, An improved procedure for mapping earthquake-induced landslide potential using a geographic information system, with applications to the Puget Sound region. p. 79-80.

McCrory, P. A., 1996, Active margin tectonics, Pacific Northwest region. p. 81-82.

Miller, K. C.; Keller, G. R., 1996, A high resolution seismic refraction and reflection survey of the Washington study corridor—Collaborative research with University of Texas at El Paso, Oregon State University, and the U.S. Geological Survey. p. 83.

Nelson, A. R.; Kelsey, H. M.; Hemphill-Haley, Eileen; Jennings, A. E., 1998, Paleoseismology of coastal Oregon. p. 15-18.

Page, R. A., 1996, USGS NEHRP fact sheets. p. 84-85.

Perkins, D. M.; Thenhaus, P. C., 1996, Reassessment of seismic sources and their earthquake recurrence rates for use in probabilistic ground motion mapping in the Pacific Northwest. p. 86-88.

Priest, G. R.; Baptista, A. M., 1998, Collaborative research between the Oregon Department of Geology and Mineral Industries and the Oregon Graduate Institute of Science and Technology—Mapping tsunami inundation hazards at Newport, Oregon. p. 28-33.

Qamar, A. I., 1996, GPS measurements of crustal deformation in the Pacific Northwest. p. 89-91.

Stephenson, W. J., 1996, 3-D boundary element modeling of the Puget Lowland thrust sheet. p. 93-94.

Thorson, R. M., 1996, Crustal motion at Lake Washington. p. 95-98.

Walsh, T. J.; Combellick, R. A.; Black, G. L., 1996, Examination of geologic effects of the 1964 Great Alaska Earthquake as a guide to recognizing paleoseismic indicators. p. 110-111.

West, M. W.; Busacca, A. J.; Berger, G. W.; Shaffer, M. E.; Ashland, F. X., 1996, A pilot study of late Quaternary surface deformation, Saddle Mountains anticline, northern Pasco Basin, Washington. p. 113.

Williams, H. F. L., 1996, Geologic record of late-Holocene earthquakes in coastal marshes—North coast and northern Puget Sound, Washington State. p. 114-115.

U.S. Geological Survey, 1997, Are walleye from Lake Roosevelt contaminated with mercury?: U.S. Geological Survey Fact Sheet 102-97, 4 p.

U.S. Geological Survey, 1998, Annual project summaries—Volume 39, Pacific Northwest: U.S. Geological Survey, 97 p. [downloaded Dec. 2, 1998, from http://erp-web.er.usgs.gov/reports/VOL39/pn/pn_vol39.htm]

Includes:

Atwater, B. F., 1998, Holocene earthquakes in western Washington. p. 1-3 [repeated p. 19-21].

Bucknam, R. C., 1998, Puget Sound paleoseismology. p. 4-11.

Campbell, N. P.; Repasky, T. R.; Ring, T. E., 1998, Earthquake hazards study in the vicinity of Toppenish Basin, south-central Washington. p. 58-60.

Crosson, R. S., 1998, Seismic velocity structure of the Puget Sound region from 3-D non-linear tomography. p. 74-78.

Frankel, A. D.; Carver, D.; Meremonte, M.; Cranswick, E.; Williams, R.; Stephenson, W. J.; Overturf, D.; Bice, T.; Odum, J. K.; Worley, D. M., 1998, Seismic hazard mapping for the Seattle region. p. 12-13.

Humphreys, E. D.; Weldon, R. J., II, 1998, Pacific Northwest deformation—Integration of regional geologic and geodetic data into kinematic and dynamic models of contemporary strain within the North American plate. p. 61-63.

Jacoby, G. C.; Bunker, D. E., 1998, Tree-ring dating of coseismic coastal subsidence in the Pacific Northwest region. p. 25-27.

Mader, G. G., 1998, Preparation of a guide to using relative earthquake hazard maps for mitigating seismic risk in the Portland metropolitan area—Collaborative research with Metro and Spangle Associates. p. 94-95.

Malone, S. D.; Crosson, R. S., 1998, Rapid strong-motion data in the Puget Sound urban area. p. 34-37.

Palmer, S. P.; Pittlekau, A. L., 1998, Liquefaction susceptibility maps for the Redmond, Kirkland, Issaquah, and Mercer Island 7.5-minute quadrangles, Washington. p. 90-93.

Priest, G. R.; Baptista, A. M., 1998, Collaborative research between the Oregon Department of Geology and Mineral Industries and the Oregon Graduate Institute of Science and Technology—Mapping tsunami inundation hazards at Newport, Oregon. p. 28-33.

Qamar, A. I., 1998, GPS investigations of Holocene fault zones in the Puget Sound basin, Washington. p. 79-83.

Qamar, A. I.; Ludwin, R. S., 1998, Database catalog of Cascadia earthquakes. p. 96-97.

Thorson, R. M., 1998, Neotectonics and shoreline change, Sammamish delta, Lake Washington. p. 71-73.

Walsh, T. J.; Logan, R. L.; Neal, K. G., 1998, Active fault investigations on the Canyon River fault, southern Olympic Range, Washington. p. 85-87.

Walsh, T. J.; Palmer, S. P.; Gerstel, W. J., 1998, Strong-motion amplification maps of the Tumwater and Lacey 1:24,000-scale quadrangles, Washington. p. 42-48.

West, M. W., 1998, A continuation of a "pilot" study of Quaternary surface deformation, Saddle Mountains anticline, northern Pasco basin, Washington. p. 49-54.

Williams, H. F. L., 1998, Geologic record of late-Holocene earthquakes in coastal marshes—North coast and northern Puget Sound, Washington State. p. 88-89.

Wong, I. G.; Bott, J. D. J.; Wright, Douglas; Sojourner, Anna; Silva, W. J.; Mabey, M. A., 1998, Deterministic and probabilistic earthquake ground shaking maps for the Portland, Oregon metropolitan area. p. 38-41.

Wagner, R. J.; Roberts, L. M., 1998, Pesticides and volatile organic compounds in surface and ground water of the Palouse subunit, central Columbia Plateau, Washington and Idaho, 1993–95: U.S. Geological Survey Water-Resources Investigations Report 97-4285, 53 p.

Wolf, S. C.; McCrory, P. A.; Hamer, M. R., 1998, Investigations of late Quaternary geologic features in Willapa Bay, Washington: U.S. Geological Survey Open-File Report 98-589, 1 sheet, scale 1:125,000.

Contract Reports

Booker, J. R., 1995, Electromagnetic remote sensing of structures controlling seismicity in the Puget Sound region: University of Washington [under contract to] U.S. Geological Survey, 9 p.

Graves, R. W., 1998, Simulating the 3D basin response in the Portland and Puget Sound regions from large subduction zone earthquakes; Final report: Woodward-Clyde Federal Services [under contract to] U.S. Geological Survey, 1 v.

Karlin, R. E., 1996?, Final report—Paleoseismicity of the Puget Sound area from geological and geophysical studies of Lake Washington: University of Nevada Mackey School of Mines [under contract to] U.S. Geological Survey, 1 v.

Malone, S. D.; Crosson, R. S.; Qamar, A. I., 1998, Cooperative operation of the Pacific Northwest Seismograph Network; Final technical report—1997: U.S. Geological Survey contract report, 1 v.

McCalpin, J. P., 1997, An improved procedure for mapping earthquake-induced landslide potential using a geographic information system, with applications to the Puget Sound region; Final technical report: GEO-HAZ Consulting, Inc. [under contract to] U.S. Geological Survey, 53 p.

Qamar, A. I., 1996, GPS measurements of crustal deformation in the Pacific Northwest: University of Washington [under contract to] U.S. Geological Survey, 6 p.

Trehu, A. M., 1997?, Collaborative research (USGS/GEOMAR/OSU)—Onshore recording of offshore shots during the 1996 USGS/GEOMAR seismic survey of the Cascadia margin; Final technical report: Oregon State University [under contract to] U.S. Geological Survey, 1 v.

West, M. W., 1997?, A continuation of a "pilot" study of Quaternary surface deformation, Saddle Mountains anticline, northern Pasco Basin, Washington—Final technical report: Michael W. West & Associates [under contract to] U.S. Geological Survey, 1 v.

OTHER REPORTS ON WASHINGTON GEOLOGY

Associated Earth Sciences, Inc., 1997, Hydrogeologic evaluation, Mats Mats quarry, Jefferson County, Washington: Associated Earth Sciences, Inc. [under contract to] Lone Star Northwest, 1 v., 4 plates.

Buchanan, J. P., 1998, Ground water and surface water hydrology of the Sullivan Road gravel pit—Comments related to the mining operation and site reclamation; Final report: John P. Buchanan [under contract to] Central Pre-Mix Concrete Company and Acme Materials and Construction Company, 46 p.

Blong, R. J., 1984, Volcanic hazards—A sourcebook on the effects of eruptions: Academic Press, 424 p.

Chelan County Public Utility District no. 1, 1976, Draft environmental impact statement for town of Rock Island vicinity, vector control project: Chelan County Public Utility District no. 1, 1 v.

Cheney, E. S., 1998, Guide to the geology of north-central Washington: Northwest Geological Society Guidebook 17, 97 p.

Clark County Regional Planning Council, 1982, Environmental impact statement, surface mining combining district and conditional use permit for rock crushing, concrete batching, asphalt mixing, Lake Development Group: Clark County Regional Planning Council, 1 v.

ENVISION Engineering Services, 1990, Draft supplemental EIS for Rand-Land surface mine and Randles Sand and Gravel—Unclassified use permit and major amendment, Frederickson area of Pierce County, WA: Pierce County Department of Planning and Natural Resource Management, 2 v.

Erickson, Denis, 1998, Yelm groundwater baseline sampling: Washington Department of Ecology Publication 98-301, 1 v.

Garrigues, R. S.; Sinclair, K. A.; Tooley, John, 1998, Chehalis river watershed surficial aquifer characterization: Washington Department of Ecology Publication 98-355, 22 p., 2 plates.

Giglio, D. F.; Erickson, Karol, 1996, Lacamas Creek watershed total maximum daily load evaluation: Washington Department of Ecology Publication 96-307, 1 v.

Golder Associates, Inc., 1998, Hydrogeologic and geomorphological study of the Selah gravel pit (near) Selah, Washington: Golder Associates, Inc., [under contract to] Central Pre-Mix Concrete Co., 1 v., 2 plates.

Hardy, John; Jacoby, Jean; Crayton, Mike, 1998, Toxic cyanobacteria—Education/lake protection; Final report: Washington Department of Health; Washington Department of Ecology Publication 98-05, 188 p.

Hartshorn, D. C.; Reidel, S. P.; Rohay, A. C., 1998, Annual Hanford seismic report for fiscal year 1998: Pacific Northwest National Laboratory PNNL-11557-9, 1 v.

Hoyle-Dodson, Guy, 1997, Puyallup Basin treatment plant metals survey: Washington Department of Ecology Publication 97-303, 1 v.

Huckell/Weinman Associates, Inc., 1991, Proposed Lakeland Hills South mining and reclamation plan and planned community development; Appendices—Draft environmental impact statement: Pierce County Department of Planning and Land Services, 1 v.

Huckell/Weinman Associates, Inc.; Associated Earth Sciences, Inc.; McCulley Frick & Gilman, Inc.; Raedeke Associates, Inc.; Art Anderson Associates; and others, 1998, Expanded environmental checklist for Northwest Aggregates Maury Island mining operation: Huckell/Weinman Associates, Inc. [under contract to Northwest Aggregates], 1 v.

Jackson, P. L.; Kimerling, A. J., editors, 1993, Atlas of the Pacific Northwest; 8th ed.: Oregon State University Press, 152 p.

Johnson, Art; Serdar, Dave; Davis, Dale, 1997, Survey for petroleum and other chemical contaminants in the sediments of Fidalgo Bay: Washington Department of Ecology Publication 97-338, 1 v.

Larson, A. G., 1996, Pesticides in Washington State's ground water—A summary report, 1988–1995: Washington Department of Ecology Publication 96-303; Washington Department of Ecology Pesticides in Ground Water 9, 22 p.

Joy, Joe; Patterson, Barbara, 1997, A suspended sediment and DDT total maximum daily load evaluation report for the Yakima River: Washington Department of Ecology Publication 97-321, 1 v.

King County Department of Development and Environmental Services, 1997, Palmer Junction gravel pit expansion, final environmental impact statement: King County Department of Development and Environmental Services, 1 v.

Komar, P. D., 1997, The Pacific Northwest coast—Living with the shores of Oregon and Washington: Duke University Press, 195 p.

Martin, J. E., editor, 1998, Contributions to the paleontology and geology of the West Coast, in honor of V. Standish Mallory: Thomas Burke Memorial Washington State Museum Research Report 6, 291 p.

Includes:

Ansfield, V. J., 1998, The Eocene Kydikabbit Point sedimentary breccias of the lower Hoko River Formation, northwestern Olympic Peninsula, Washington, and their contained invertebrates. p. 1-21.

McDougall, K. A., 1998, Late Eocene neritic and bathyal foraminiferal assemblages of the Pacific Northwest. p. 43-60.

Richardson, S. L., 1998, West Coast Tertiary foraminiferal literature—A selected bibliography. p. 73-90.

Wehr, W. C., 1998, Middle Eocene insects and plants of the Okanogan Highlands. p. 99-109.

Pelletier, G. J., 1997, Colville River water quality—Pollutant loading capacity and recommendations for total maximum daily loads: Washington Department of Ecology Publication 96-349, 1 v.

Pelto, M. S., 1997, Changes in glaciers on Glacier Peak in the last 100 years: [Privately published by the author, 8 p.]

Pringle, P. T.; Hickson, C. J., 1998, Cascade volcanoes—Processes and hazards: International Association for Engineering Geology and the Environment, 8th Congress, Technical Tour Guide Book Trip 14, 22 p.

Raymond, J. R., 1955, Applicability of geophysical methods of exploration at Hanford Atomic Products Operation: Hanford Atomic Products Operation HW-38707, 14 p.

Rector, Julie; Hallock, Dave, 1995, 1994 statewide water quality assessment lakes chapter—Companion document to Washington State's 305(b) report: Washington Department of Ecology Publication 95-311, 60 p.

Schlichting, Robert; Peterson, C. D., 1998, A reconnaissance of freshwater marsh stratigraphy for evidence of tsunami-induced catastrophic marine flooding, Grayland and Long Beach, Washington: Portland State University [under contract to] Washington Department of Natural Resources, 1 v.

Sinclair, K. A., 1998, Basic American Foods class II groundwater inspection: Washington Department of Ecology Publication 98-330, 1 v.

Sinclair, Kelsey; Bentley, R. D., 1998, Stratigraphic correlation of the N_2 Grande Ronde basalt across the Kittitas Valley, Washington State: Central Washington University Department of Geology, 1 v., 1 plate.

Smith, A. K.; Rector, Julie, 1997, Water quality assessments of selected lakes within Washington State, 1994: Washington Department of Ecology Publication 97-307, 1 v

Stewart, R. J., 1998, Field trip to the western Olympic Mountains, Washington—An excursion into the geology of the Olympic subduction complex, guided by the mapping of Rowland W. Tabor, Weldon W. Rau, Parke D. Snavely, Jr., and Wallace M. Cady, with observations (and obfuscations), by R. J. Stewart: Northwest Geological Society Guidebook 16, 1 v.

Stroud, W. P.; Droullard, R. F., 1996, 1995 Midnite mine radiation survey: U.S. Bureau of Mines Report of Investigations 9610, 25 p.

Thorsen, G. W.; Shipman, Hugh, 1998, Bluff erosion monitoring on Puget Sound—A guide for volunteers: Washington Department of Ecology [under contract to] Island County Beach Walkers, 79 p.

Torres, R. L.; Luehring, R. W., 1996, Corrective action study, geotechnical issues at Salmon Lake dam: U.S. Bureau of Reclamation Technical Memorandum OZ-8313-5, 1 v.

U.S. Army Corps of Engineers, 1997, Post event report—Winter storm of 1996–97, Federal Disaster DR 1159, western Washington summary; Final document 16 May 1997: U.S. Federal Emergency Management Agency, 1 v.

U.S. Environmental Protection Agency, 1998, Mineralogical study of borehole MW-206, ASARCO smelter site, Tacoma, Washington: U.S. Environmental Protection Agency, 1 v.

U.S. Federal Emergency Management Agency, 1996, Interagency Hazard Mitigation Team report, including progress report on early implementation strategies—State of Washington, winter storms of 1995–1996; FEMA-DR-1079, declared January 3, 1996; FEMA-DR-1100-WA, declared February 9, 1996: U.S. Federal Emergency Management Agency, 88 p.

U.S. Federal Emergency Management Agency, 1997, Hazard Mitigation Survey Team report for FEMA DR-1152-WA, declared January 7, 1997 and FEMA DR-1159-WA, declared January 17, 1997, including addendum report for FEMA DR-1172-WA, declared April 2, 1997: U.S. Federal Emergency Management Agency, 123 p.

U.S. Federal Emergency Management Agency, 1997, Supplemental flood hazard mitigation report, in response to the November 26, 1990 disaster declaration, State of Washington, FEMA-883-DR-WA; A supplemental report covering—San Juan, Kitsap, Pierce, Thurston, Lewis, Grays Harbor Pacific, Wahkiakum, Mason, Chelan, Yakima, Kittitas, Island, Jefferson Counties: U.S. Federal Emergency Management Agency, 44 p.

U.S. Federal Emergency Management Agency, 1998, The Project Impact hazard mitigation guidebook for northwest communities—Alaska, Idaho, Oregon, Washington: U.S. Federal Emergency Management Agency, 58 p.

U.S. National Geophysical Data Center, 1997, Geologic hazards photos—Vol. 1, Earthquakes, general topics; Vol. 2, Earthquake events; Vol. 3, Landslides, tsunamis, and volcanoes: U.S. National Geophysical Data Center, 3 compact disks.

U.S. National Weather Service, 1997, Disastrous floods from the severe winter storms in California, Nevada, Washington, Oregon, and Idaho—December 1996–January 1997: U.S. National Oceanic and Atmospheric Administration Natural Disaster Survey Report, 1 v.

U.S. National Oceanic and Atmospheric Administration, 1998, NOAA's estuarine eutrophication survey; Volume 5—Pacific Coast region: U.S. National Oceanic and Atmospheric Administration, 75 p.

University of Washington Geophysics Program, 1998, Quarterly network report 97-D on seismicity of Washington and Oregon: University of Washington Geophysics Program, 34 p.

University of Washington Geophysics Program, 1998, Quarterly network report 98-B on seismicity of Washington and Oregon April 1 through June 30, 1998: University of Washington Geophysics Program, 32 p.

Washington Department of Natural Resources, 1997, Lake Whatcom watershed analysis: Washington Department of Natural Resources, 1 v.

Washington Department of Natural Resources, 1998, Our changing nature—Natural resource trends in Washington State: Washington Department of Natural Resources, 75 p.

Washington Department of Transportation, 1998, SR16/Union Avenue vicinity to SR302 vicinity, draft environmental impact statement summary: Washington Department of Transportation, 1 v. [downloaded 10/23/98 from http://www.wsdot.wa.gov/solve16/deis.html]

PAPERS ON WASHINGTON GEOLOGY

Aalto, K. R.; Sharp, W. D.; Renne, P. R., 1998, $^{40}Ar/^{39}Ar$ dating of detrital micas from Oligocene–Pleistocene sandstones of the Olympic Peninsula, Klamath Mountains, and northern California coast ranges—Provenance and paleodrainage patterns: Canadian Journal of Earth Sciences, v. 35, no. 7, p. 735-745.

Alidibirov, A.; Dingwell, D. B.; Stevenson, R. J.; Hess, K.-U.; Webb, S. L.; Zinke, J., 1997, Physical properties of the 1980 Mount St. Helens cryptodome magma: Bulletin of Volcanology, v. 59, no. 2, p. 103-111.

Anderson, C. H., Jr.; Behrens, C. J.; Floyd, G. A.; Vining, M. R., 1998, Crater firn caves of Mount St. Helens, Washington: Journal of Cave and Karst Studies, v. 60, no. 1, p. 44-50.

Anderson, R. T.; Chapelle, F. H.; Lovley, D. R., 1998, Evidence against hydrogen-based microbial ecosystems in basalt aquifers: Science, v. 281, no. 5379, p. 976-977.

Anderson, S. W.; Stofan, E. R.; Plaut, J. J.; Crown, D. A., 1998, Block size distributions on silicic lava flow surfaces—Implications for emplacement conditions: Geological Society of America Bulletin, v. 110, no. 10, p. 1258-1267.

Baffaut, C.; Nearing, M. A.; Nicks, A. D., 1996, Impact of CLIGEN parameters on WEPP-predicted average annual soil loss: American Society of Agricultural Engineers Transactions, v. 39, no. 2, p. 447-457.

Baker, E. T.; Massoth, G. J.; Feely, R. A.; Cannon, G. A., 1998, The rise and fall of the CoAxial hydrothermal site, 1993–1996: Journal of Geophysical Research, v. 103, no. B5, p. 9791-9806.

Barendregt, R. W.; Irving, Edward, 1998, Changes in the extent of North American ice sheets during the late Cenozoic: Canadian Journal of Earth Sciences, v. 35, no. 5, p. 504-509.

Barnes, S. M.; Nierzwicki-Bauer, S. A., 1997, Microbial diversity in ocean, surface and subsurface environments. *In* Banfield, J. F.; Nealson, K. H., editors, Geomicrobiology—Interactions between microbes and minerals: Mineralogical Society of America Reviews in Mineralogy, v. 35, p. 35-79.

Belousov, A. B.; Belousova, M. G., 1997, Gigantskiye obvaly na vulkanakh [Gigantic landslides on volcanoes]: Priroda, v. 11, p. 70-81. (In Russian)

Bottari, A.; Caccamo, D.; Malone, S. D.; Neri, G., 1987, Statistical analysis of Mt. St. Helens earthquakes. *In* Consiglio Nazionale delle Ricerche Gruppo Nazionale per la Vulcanologia, Bollettino 1987: Consiglio Nazionale delle Ricerche, p. 145-157.

Bradford, J. H.; Sawyer, D. S.; Zelt, C. A.; Oldow, J. S., 1998, Imaging a shallow aquifer in temperate glacial sediments using seismic reflection profiling with DMO processing: Geophysics, v. 63, no. 4, p. 1248-1256.

Brandon, A. D.; Draper, D. S., 1998, Reply to the comment by B. R. Frost and C. Ballhaus on "Constraints on the origin of the oxidation state of mantle overlying subduction zones—An example from Simcoe, Washington, USA": Geochimica et Cosmochimica Acta, v. 62, no. 2, p. 333-335.

Brandon, M. T.; Roden-Tice, M. K.; Garver, J. I., 1998, Late Cenozoic exhumation of the Cascadia accretionary wedge in the Olympic Mountains, northwest Washington State: Geological Society of America Bulletin, v. 110, no. 8, p. 985-1009; GSA Data Repository item 9865.

Braun, Jill; Major, Graeme; West, D. O.; Bukovansky, Michal, 1998, Geologic hazards evaluation boosts risk-management program for western U.S. pipeline: Oil and Gas Journal, v. 96, no. 45, p. 73-79.

Cameron, D. E., 1995, Structural setting and features of Au-Ag orebodies at the Cannon mine, Wenatchee, Washington. *In* Coyner, A. R.; Fahey, P. L., editors, Geology and ore deposits of the American Cordillera; Symposium proceedings: Geological Society of Nevada, v. 2, p. 1089-1110.

Chadwick, W. W., Jr.; Embley, R. W., 1998, Graben formation associated with recent dike intrusions and volcanic eruptions on the mid-ocean ridge: Journal of Geophysical Research, v. 103, no. B5, p. 9807-9825.

Cheney, E. S.; Trammell, J. W., 1995, The "inside-out" Quartz Creek quartz dioritic, batholithic, copper deposit, central Cascade Range, Washington. *In* Coyner, A. R.; Fahey, P. L., editors, Geology and ore deposits of the American Cordillera; Symposium proceedings: Geological Society of Nevada, v. 3, p. 1521-1532.

Colpron, Maurice; Warren, M. J.; Price, R. A., 1998, Selkirk fan structure, southeastern Canadian Cordillera—Tectonic wedging against an inherited basement ramp: Geological Society of America Bulletin, v. 110, no. 8, p. 1060-1074.

Conca, J. L.; Wright, Judith, 1998, The UFA method for rapid, direct measurements of unsaturated transport properties in soil, sediment, and rock: Australian Journal of Soil Research, v. 36, no. 2, p. 291-315.

Davis, E. E.; Becker, Keir, 1998, Borehole observatories record driving forces for hydrothermal circulation in young oceanic crust: Eos (American Geophysical Union Transactions), v. 79, no. 31, p. 369, 377-378.

Endo, E. T.; Murray, T. L.; Power, J. A., 1996, A comparison of pre-eruption real-time seismic amplitude measurements for eruptions at Mount St. Helens, Redoubt Volcano, Mount Spurr, and Mount Pinatubo. *In* Newhall, C. G.; Runongbayan, R. S., editors, Fire and mud—Eruptions and lahars of Mount Pinatubo, Philippines: Philippine Institute of Volcanology and Seismology; University of Washington Press, p. 233-247.

Engineering & Mining Journal, 1998, Vulcan software system—For exploration and mine design, it's worth its weight in gold: Engineering & Mining Journal, v. 199, no. 7, p. 32NN-32PP.

Evans, R. L.; Webb, S. C.; Jegen, Marion; Sananikone, Khamla, 1998, Hydrothermal circulation at the Cleft–Vance overlapping spreading center—Results of a magnetometric resistivity survey: Journal of Geophysical Research, v. 103, no. B6, p. 12,321-12,338.

Field, C. W., 1993, Minerals and mining. *In* Jackson, P. L.; Kimerling, A. J., editors, Atlas of the Pacific Northwest; 8th ed.: Oregon State University Press, p. 122-127.

Fifarek, R. H.; Devlin, B. D.; Tschauder, R. J., Jr., 1995, Au-Ag mineralization at the Golden Promise deposit, Republic district, Washington—Relation to graben development and hot springs processes. *In* Coyner, A. R.; Fahey, P. L., editors, Geology and ore deposits of the American Cordillera; Symposium proceedings: Geological Society of Nevada, v. 2, p. 1063-1088.

Fisk, M. R.; Giovannoni, S. J.; Thorseth, I. H., 1998, Alteration of oceanic volcanic glass—Textural evidence of microbial activity: Science, v. 281, no. 5379, p. 978-980.

Fountain, A. G.; Jacobel, R. W., 1997, Advances in ice radar studies of a temperate alpine glacier, South Cascade Glacier, Washington, U.S.A.: Annals of Glaciology, v. 24, p. 303-308.

Frost, B. R.; Ballhaus, Chris, 1998, Comment on "Constraints on the origin of the oxidation state of mantle overlying subduction zones—An example from Simcoe, Washington, USA," by A. D. Brandon and D. S. Draper: Geochimica et Cosmochimica Acta, v. 62, no. 2, p. 329-331.

Glicken, Harry, 1998, Rockslide-debris avalanche of May 18, 1980, Mount St. Helens volcano, Washington: Geological Survey of Japan Bulletin, v. 49, no. 2/3, p. 55-106.

Hachey, J. E.; Plum, R. L.; Byrne, R. J.; Kilian, A. P.; Jenkins, D. V., 1994, Blast densification of a thick, loose debris flow at Mt. St. Helens, Washington. *In* Yeung, A. T.; Felio, G. Y., editors, Vertical and horizontal deformations of foundations and embankments: American Society of Civil Engineers Geotechnical Special Publication 40, v. 1, p. 502-512.

Halbert, W. E.; Jensen, R. E., 1996, Influence of tidal fluctuations on coastal aquifers—General principles and case studies. *In* Proceedings of the Tenth National Outdoor Action Conference and Exposition—Aquifer Remediation/Ground Water Monitoring/Geophysical Methods: National Ground Water Association, p. 575-592.

Harris, C. F. T., 1998, Washington State's 1:100,000-scale geologic map database—An Arc/Info data model example. *In* Soller, D. R., editor, Digital mapping techniques '98—Workshop proceedings: U.S. Geological Survey Open-File Report 98-487, p. 27-35.

Harris, D. M.; Rose, W. I., 1996, Dynamics of carbon dioxide emissions, crystallization, and magma ascent—Hypotheses, theory, and applications to volcano monitoring at Mount St. Helens: Bulletin of Volcanology, v. 58, no. 2-3, p. 163-174.

Hartnett, H. E.; Keil, R. G.; Hedges, J. I.; Devol, A. H., 1998, Influence of oxygen exposure time on organic carbon preservation in continental margin sediments: Nature, v. 391, no. 6667, p. 572-574.

Iverson, R. M.; Schilling, S. P.; Vallance, J. W., 1998, Objective delineation of lahar-inundation hazard zones: Geological Society of America Bulletin, v. 110, no. 8, p. 972-984.

Kalbasi, M.; Peryea, F. J.; Lindsay, W. L.; Drake, S. R., 1995, Measurement of divalent lead activity in lead arsenate contaminated soils: Soil Science Society of America Journal, v. 59, no. 5, p. 1274-1280.

Keil, R. G.; Montluçon, D. B.; Prahl, F. G.; Hedges, J. I., 1994, Sorptive preservation of labile organic matter in marine sediments: Nature, v. 370, no. 6490, p. 549-552.

Keil, R. G.; Tsamakis, Elizabeth; Giddings, J. C.; Hedges, J. I., 1998, Biochemical distributions (amino acids, neutral sugars, and lignin phenols) among size-classes of modern marine sediments from the Washington coast: Geochimica et Cosmochimica Acta, v. 62, no. 8, p. 1347-1364.

Khire, M. V.; Benson, C. H.; Bosscher, P. J., 1997, Water balance modeling of earthen final covers: Journal of Geotechnical and Geoenvironmental Engineering, v. 123, no. 8, p. 744-754.

Klotz, Sally; Smith, Daphne, 1975, Distribution of plant species on the mounds of Mima Prairie, Thurston County, Washington. *In* Herman, S. G.; Wiedemann, A. M., editors, Contributions to the natural history of the southern Puget Sound region, Washington: The Evergreen State College, p. 25-62.

Knitter, C. C.; Haskell, K. G.; Peterson, M. L., 1993, Use of low plasticity silt for soil liners and covers. *In* Prakash, Shamsher, editor, Third International Conference on Case Histories in Geotechnical Engineering: University of Missouri–Rolla, v. II, p. 1255-1259.

Koler, T. E., 1998, Evaluating slope stability in forest uplands with deterministic and probabilistic models: Environmental and Engineering Geoscience, v. 4, no. 2, p. 185-194.

Kuuskraa, V. A.; Schmoker, J. W.; Dyman, T. S., 1998, Emerging U.S. gas resources—6 (conclusion); Diverse gas plays lurk in gas resource pyramid: Oil & Gas Journal, v. 96, no. 23, p. 123-130.

Lehrsch, G. A., 1998, Freeze-thaw cycles increase near-surface aggregate stability: Soil Science, v. 163, no. 1, p. 63-70.

Lewis, D. E.; Comis, J. G.; Prych, E. A., 1987, Flood protection in the Puyallup River Basin, Washington—Past, present and future. *In* Association of State Floodplain Managers, Realistic approaches to better floodplain management; 11th annual conference, Proceedings: University of Colorado Natural Hazards Information Center Special Publication 18, p. 298-304.

Long, A. J.; Shennan, Ian, 1998, Models of rapid relative sea-level change in Washington and Oregon, USA: The Holocene, v. 8, no. 2, p. 129-142.

Lowe, C. H.; Larson, P. B., 1995, Geology of the Key East gold deposit, Ferry County, Washington. *In* Coyner, A. R.; Fahey, P. L., editors, Geology and ore deposits of the American Cordillera; Symposium proceedings: Geological Society of Nevada, v. 2, p. 1111-1132.

Lyatsky, H. V., 1997, Control of the western continental margin of North America by the Cordilleran Fairweather–Queen Charlotte–Wallowa fault system. *In* Cook, F.; Erdmer, P., compilers, Slave–Northern Cordillera Lithospheric Evolution (SNORCLE) Transect and Cordilleran Tectonics Workshop Meeting (March 7–9): University of Calgary Lithoprobe Report 56, p. 184-186.

McGee, K. A.; Sutton, A. Jefferson, 1994, Eruptive activity at Mount St. Helens, Washington, USA, 1984–1988—A gas geochemistry perspective: Bulletin of Volcanology, v. 56, no. 6-7, p. 435-446.

Monastersky, Richard, 1998, Waves of death: Science News Online, 10/3/98, 7 p. [downloaded 10/21/98 from http://www.sciencenews.org/sn_arc98/10_3_98/Bob2.htm].

Murphy, J. B.; Oppliger, G. L.; Brimhall, G. H., Jr.; Hynes, Andrew, 1998, Plume-modified orogeny—An example from the western United States: Geology, v. 26, no. 8, p. 731-734.

Mustard, P. S.; Clague, J. J.; Woodsworth, G. J.; Hickson, C. J.; Jackson, L. E., Jr.; Luternauer, J. L.; Monger, J. W. H.; Richetts, B. D.; and others, 1998, Geology and geological hazards of the greater Vancouver area. *In* Karrow, P. F.; White, O. l., editors, Urban geology of Canadian cities: Geological Association of Canada Special Paper 42, p. 39-70.

Myagkov, N. N., 1998, Model of a strong volcanic blast and a method of estimating the mass ejected: Geophysical Journal International, v. 133, no. 1, p. 209-211.

O'Connor, J. E.; Costa, J. E., 1993, Geologic and hydrologic hazards in glacierized basins in North America resulting from 19th and 20th century global warming: Natural Hazards, v. 8, no. 2, p. 121-140.

Orange, D. L.; Knittle, Elise; Farber, Daniel; Williams, Quentin, 1996, Raman spectroscopy of crude oils and hydrocarbon fluid inclusions—A feasibility study. *In* Dyar, M. D.; McCannon, Catherine; Schaefer, M. W., editors, Mineral spectroscopy—A tribute to Roger G. Burns: Geochemical Society Special Publication 5, p. 65-81.

Paterson, S. R.; Miller, R. B., 1998, Magma emplacement during arc-perpendicular shortening—An example from the Cascades crystalline core, Washington: Tectonics, v. 17, no. 4, p. 571-586.

Pearce, T. H., 1994, Recent work on oscillatory zoning in plagioclase. *In* Parsons, Ian, editor, Feldspars and their reactions: Kluwer Academic Publishers NATO ASI series C, v. 421, p. 313-349.

Pierce, J. K., 1998, The big one: Seattle Magazine, v. 7, no. 10, p. 30-35.

Pierson, T. C., 1998, An empirical method for estimating travel times for wet volcanic mass flows: Bulletin of Volcanology, v. 60, no. 2, p. 98-109.

Prahl, F. G.; Small, L. F.; Sullivan, B. A.; Cordell, J.; Simenstad, C. A.; Crump, B. C.; Baross, J. A., 1998, Biogeochemical gradients in the lower Columbia River: Hydrobiologia, v. 361, p. 37-52.

Rabinowicz, M.; Boulegue, J.; Genthon, P., 1998, Two- and three-dimensional modeling of hydrothermal convection in the sedimented Middle Valley segment, Juan de Fuca Ridge: Journal of Geophysical Research, v. 103, no. B10, p. 24,045-24,065.

Ray, C. E.; Domning, D. P.; McKenna, M. C., 1994, A new specimen of *Behemotops proteus* (order Desmostylia) from the marine Oligocene of Washington. *In* Berta, Annalisa; Demere, T. A., editors, Contributions in marine mammal paleontology honoring Frank C. Whitmore, Jr.: San Diego Society of Natural History Proceedings 29, p. 205-222.

Reidel, S. P., 1998, Emplacement of Columbia River flood basalt: Journal of Geophysical Research, v. 103, no. B11, p. 27,393-27,410.

Reuter, R. J.; McDaniel, P. A.; Hammel, J. E.; Falen, A. L., 1998, Solute transport in seasonal perched water tables in loess-derived soilscapes: Soil Science Society of American Journal, v. 62, no. 4, p. 977-983.

Riedell, K. B.; Jones, B. K.; Cheney, E. S., 1995, Geology, geochemistry, and evolution of the Mazama porphyry copper-molybdenum-gold deposit, Okanogan County, Washington. *In* Coyner, A. R.; Fahey, P. L., editors, Geology and ore deposits of the American Cordillera; Symposium proceedings: Geological Society of Nevada, v. 3, p. 1501-1520.

Rosenfeld, C. L., 1993, Landforms and geology. *In* Jackson, P. L.; Kimerling, A. J., editors, Atlas of the Pacific Northwest; 8th ed.: Oregon State University Press, p. 40-47.

Rychkun, Ed, 1995, 3D geostatistical modeling applied to radioactive waste site: Earth Observation Magazine, v. 4, no. 10, p. 42-44.

Schilling, S. P.; Iverson, R. M., 1997, Automated, reproducible delineation of zones at risk from inundation by large volcanic debris flows. *In* Chen, C.-L., editor, Debris-flow hazards mitigation—Mechanics, prediction, and assessment: American Society of Civil Engineers, p. 176-186.

Sohn, R. A.; Hildebrand, J. A.; Webb, S. C., 1998, Postrifting seismicity and a model for the 1993 diking event on the CoAxial segment, Juan de Fuca Ridge: Journal of Geophysical Research, v. 103, no. B5, p. 9867-9877.

Stein, J. S.; Fisher, A. T.; Langseth, M. G.; Jin, W.; Iturrino, G. J.; Davis, E. E., 1998, Fine-scale heat flow, shallow heat sources, and decoupled circulation systems at two sea-floor hydrothermal sites, Middle Valley, northern Juan de Fuca Ridge: Geology, v. 26, no. 12, p. 1115-1118.

Tedford, R. H.; Barnes, L. G.; Ray, C. E., 1994, The early Miocene littoral ursoid carnivoran *Kolponomos*—Systematics and mode of life. *In* Berta, Annalisa; Demere, T. A., editors, Contributions in marine mammal paleontology honoring Frank C. Whitmore, Jr.: San Diego Society of Natural History Proceedings 29, p. 11-32.

Tesoriero, A. J.; Voss, F. D., 1997, Predicting the probability of elevated nitrate concentrations in the Puget Sound basin—Implications for aquifer susceptibility and vulnerability: Ground Water, v. 35, no. 6, p. 1029-1039.

Testa, S. M., 1996, Considerations and methodology in using natural attenuation as a remedial strategy. *In* Proceedings of the Tenth National Outdoor Action Conference and Exposition—Aquifer remediation/ground water monitoring/geophysical methods: National Ground Water Association, p. 283-302.

Thimsen, C. A.; Keil, R. G., 1998, Potential interactions between sedimentary dissolved organic matter and mineral surfaces: Marine Chemistry, v. 62, no. 1-2, p. 65-76.

Thordarson, Th.; Self, Stephen, 1998, The Roza Member, Columbia River Basalt Group—A gigantic pahoehoe lava flow field formed by endogenous processes?: Journal of Geophysical Research, v. 103, no. B11, p. 27,411-27,445.

Thornton, P. E.; Running, S. W.; White, M. A., 1997, Generating surfaces of daily meteorological variables over large regions of complex terrain: Journal of Hydrology, v. 190, no. 3-4, p. 214-251.

Verardo, D. J., 1998, A river runs through it: GSA Today, v. 8, no. 8, p. 12-15.

Wells, R. E.; Weaver, C. S.; Blakely, R. J., 1998, Fore-arc migration in Cascadia and its neotectonic significance: Geology, v. 26, no. 8, p. 759-762.

Wigmosta, M. S.; Burges, S. J., 1997, An adaptive modeling and monitoring approach to describe the hydrologic behavior of small catchments: Journal of Hydrology, v. 202, nos. 1-4, p. 48-77.

Wold, W. L., 1996, GIS and watershed analysis—Not just for mapping anymore! *In* Hallam, C. A.; Salisbury, J. M.; Lanfear, K. J.; Battaglin, W. A., editors, Proceedings of the AWRA Annual Symposium, GIS and Water Resources: American Water Resources Association Technical Publication Series TPS-96-3, p. 317-326.

Wood, A. W.; Lettenmaier, D. P.; Palmer, R. N., 1997, Assessing climate change implications for water resources planning: Climate Change, v. 37, no. 1, p. 203-228.

Zabowski, Darlene; Everett, R. L., 1997, Extractable metals and plant uptake with amelioration and revegetation of abandoned copper mine tailings. *In* Iskandar, I. K.; Adriano, D. C., editors, Remediation of soils contaminated with metals: Science Reviews Advances in Environmental Science, p. 111-122.

OTHER REPORTS OF INTEREST

Applied Technology Council, 1997, Seismic design criteria for bridges and other highway structures—Current and future: Applied Technology Council ATC-18, 152 p.

Baker, Ron, 1996, A primer of oilwell drilling—A basic text of oil and gas drilling; 5th ed. (rev.): University of Texas at Austin Petroleum Extension Service, 184 p., 1 plate.

Barksdale, R. D., editor, 1991, The aggregate handbook: National Stone Association, 1 v.

Boland, M. A.; Piccione, C. L., editors, 1998, Earth-science education resource directory; 2nd ed.: American Geological Institute, 270 p.

Booth, D. B.; Collins, B. D.; Montgomery, D. R.; Perkins, S. J., 1997, Geology and geomorphology of stream channels—Course manual: University of Washington Engineering Professional Programs, 1 v.

Brodzikowski, K.; van Loon, A. J., 1991, Glacigenic sediments: Elsevier Developments in Sedimentology 49, 674 p.

Dmowska, Renata; Saltzman, Barry, editors, 1999, Tsunamigenic earthquakes and their consequences: Academic Press Advances in Geophysics, v. 39, 215 p.

Ferris, F. K.; Kleinman, L. H.; and others, 1996, Handbook of western reclamation techniques: U.S. Office of Surface Mining Reclamation and Enforcement, 1 v.

Huzziff, C. A.; Ronan, K. R., 1998, Prediction of children's coping following a natural disaster—The Mount Ruapehu eruptions—A prospective study: Journal of Traumatic Stress [submitted manuscript, under review July 1998], 38 p.

Johnson, D. M.; Hooper, P. R.; Conrey, R. M., 1998, XRF analysis of rocks and minerals for major and trace elements on a single low dilution Li-tetraborate fused bead: Washington State University GeoAnalytical Laboratory, 25 p.

King County Department of Natural Resources Water and Land Resources Division, 1994, Northwest native plants—Identification and propagation for revegetation and restoration projects: King County Department of Natural Resources, 1 v.

Laprade, W. T., 1998, The sculptor and the geologist—Turning stone into art: Geotimes, v. 43, no. 7, p. 20-23.

O'Connor, J. E.; Burns, Scott, 1998, Beyond the Channeled Scablands—Quaternary geology of the Willamette Valley, Oregon, September 19 & 20, 1998: Friends of the Pleistocene Pacific Northwest Cell, 6th Annual Meeting, Field Trip, 72 p.

Palmer, Gayle; Stevenson, Shanna, editors, 1992, Thurston County place names—A heritage guide: Thurston County Historic Commission, 140 p., 1 plate.

Priest, G. R.; Myers, E. P., III; Baptista, A. M.; Kamphaus, R. A.; Peterson, C. D.; Darienzo, M. E., 1997, Tsunami hazard map of the Yaquina Bay area, Lincoln County, Oregon: Oregon Department of Geology and Mineral Industries Interpretive Map Series IMS-2, 1 sheet, scale 1:12,000.

Ronan, K. R., 1997, The effects of a "benign" disaster—Symptoms of post-traumatic stress in children following a series of volcanic eruptions: Australasian Journal of Disaster and Trauma Studies, 1997/1 [downloaded Oct. 12, 1998, from http://www.massey.ac.nz/~trauma/issues/1997-1/ronan.htm], 9 p.

Ronan, K. R.; Johnston, D. M., 1997, Children's risk perceptions and preparedness—A hazards education survey: Auckland City Council; Auckland Regional Council, 1 v.

Ronan, K. R.; Johnston, D. M., 1998, The effects of time and brief school-based intervention on self-reported symptoms of PTSD and coping following a series of volcanic eruptions: Journal of Consulting and Clinical Psychology [submitted manuscript, under review July 1998], 19 p.

Teller, A. E., 1994, Fundamentals of blasting and reclamation blasting workshop: Washington Division of Geology and Earth Resources unpublished report, 1 v.

Wilson, W. E.; Moore, J. E., editors, 1998, Glossary of hydrology: American Geological Institute, 248 p. ■

HOW TO FIND OUR MAIN OFFICE



Washington Bibliography Available on CD-ROM

The *Digital Index to the Geology and Mineral Resources of Washington, 1798 through July 1998,* compiled and edited by Connie J. Manson, is now available on CD-ROM. The file contains the citations and indexing for more than 32,000 items and includes both the items listed in our printed bibliographies and those non-Washington items held in our library. The disk contains the search software and runs on Windows 3.1 or higher. It sells for $3.22 + .28 tax (for Washington residents only) = $3.50. (Please include $1.00 postage and handling for each order.)

DIVISION PUBLICATIONS

New Releases

Quaternary Stratigraphy, Cross Sections, and General Geohydrologic Potential of the Bow and Alger 7.5-minute Quadrangles, Western Skagit County, Washington, Open File Report 98-8, by Joe D. Dragovich and Carly L. Grisamer. This report with 30 p. text and 6 plates is based on interpretations of numerous well logs. $5.56 + .44 tax = $6.00.

Geologic and Geophysical Mapping of Washington, 1984 through 1999, and Theses on the Geology of Washington, 1986 through 1998, Open File Report 99-1, compiled by C. J. Manson. 55 pages, 9 plates. Listed are 264 geologic or geophysical maps and 508 theses. This report supersedes (by updating) Open File Report 98-1. $1.39 + .11 = $1.50.

1999 Gold and Fish—Rules and Regulations for Mineral Prospecting and Placer Mining in Washington State has just come in. It will be available free of charge.

Orders must be prepaid. Make check or money order payable to the Department of Natural Resources. Taxes apply to Washington residents only. Please include $1.00 for postage and handling of orders to be sent by mail.

Northwest Geological Society Website

The Northwest Geological Society now has a website at http://www.scn.org/tech/nwgs/. Much of the information on the website is not specifically about NWGS—it focuses on Northwest geology in general. Editor/webmaster Dave Knoblach is trying to improve communication between NW geoscientists and geology-related organizations. "Since the website went on line in September 1998, I have found several organizations previously unknown to me from Vancouver, B.C., to Portland to Spokane. I am now in regular communication with most of them. The site has received attention from the Tri-Cities, Ellensburg, Moses Lake, and Walla Walla, judging from the e-mail I have received and links that I have found to our site from other organizations. A number of non-NWGS organizations and individuals have started to contribute to the NWGS website," he said.

The NWGS website contains a robust calendar page from many organizations in the northwest, geology link pages, and links to government and educational agencies, weather, employment, and professional organizations. It also contains announcements focused mainly on Washington geology and the NWGS. ■

APOLOGIES TO OUR READERS

Due to loss of staff and an increased work load, *Washington Geology* has been behind schedule for most of the last year. We hope to be back on track soon, although we may have to cut back on the size and (or) frequency of the issues. Thank you for your patience.

Weldon Rau Honored

Weldon W. Rau, Division of Geology and Earth Resource geologist emeritus, has been honored with a mention in *Oregon Fossils* (Orr and Orr, in press). "Weldon Rau, who performed much of the biostratigraphic micropaleontology work in the Pacific Northwest from the middle 1960s onward, enjoyed a reputation as a rapid but careful worker. Born in Tacoma in 1921, Rau completed his graduate education at the University of Iowa where he earned a Ph.D. in paleontology in 1950. Taking a job as a micropaleontologist and stratigrapher with the Fuels Branch of the U.S. Geological Survey that same year, he specialized in benthic (bottom-dwelling) Tertiary foraminifera of the Pacific Northwest. Several extensive monographs on the Olympic Peninsula in addition to descriptions of many isolated faunas and local geology were the result. Two of his most interesting bulletins are on the scenic geology and history of the Washington coast (Rau, 1973, 1980)....Weldon joined the Washington Division of Geology and Earth Resources in 1960 and since retirement continues to work there at his office in Olympia."

A large gastropod, *Abyssochrysos raui* Goedert & Kaler, sp. nov., was named for Rau in recognition of his pioneering work on the foraminiferal biostratigraphy of Tertiary marine rocks in the Pacific Northwest by Goedert and Kaler (1996).

References

Goedert, J. L.; Kaler, K. L., 1996, A new species of *Abyssochrysos* (Gastropoda: Loxonematoidea) from a middle Eocene cold-seep carbonate in the Humptulips Formation, Western Washington: The Verliger, v. 39, no. 1, p. 65-70.

Orr, E. L; Orr W. N., in press, Oregon fossils: Kendall/Hunt.

Rau, W. W., 1973, Geology of the Washington coast between Point Grenville and the Hoh River: Washington Division of Geology and Earth Resources Bulletin 66, 58 p., 72 figs. [color, $3.00]

Rau, W. W., 1980, Washington coastal geology between the Hoh and Quillayute Rivers: Washington Division of Geology and Earth Resources Bulletin 72, 57 p., 74 figs. [color, $5.00] ■

WASHINGTON STATE DEPARTMENT OF
Natural Resources
Jennifer M. Belcher - Commissioner of Public Lands

Department of Natural Resources
Division of Geology and Earth Resources
PO Box 47007
Olympia, WA 98504-7007

BULK RATE
U.S. POSTAGE PAID
Washington State
Department of Printing

ADDRESS SERVICE REQUESTED